

BLOCK-PROXIMAL METHODS WITH SPATIALLY ADAPTED ACCELERATION

Tuomo Valkonen*

V4: 2018-11-19[†]

Abstract We study and develop (stochastic) primal–dual block-coordinate descent methods for convex problems based on the method due to Chambolle and Pock. Our methods have known convergence rates for the iterates and the ergodic gap: $O(1/N^2)$ if each block is strongly convex, $O(1/N)$ if no convexity is present, and more generally a mixed rate $O(1/N^2) + O(1/N)$ for strongly convex blocks, if only some blocks are strongly convex. Additional novelties of our methods include blockwise-adapted step lengths and acceleration, as well as the ability to update both the primal and dual variables randomly in blocks under a very light compatibility condition. In other words, these variants of our methods are doubly-stochastic. We test the proposed methods on various image processing problems, where we employ pixelwise-adapted acceleration.

1 INTRODUCTION

We want to efficiently solve optimisation problems of the form

$$(P_0) \quad \min_x G(x) + F(Kx),$$

arising, in particular, from image processing and inverse problems. We assume $G : X \rightarrow \overline{\mathbb{R}}$ and $F : Y \rightarrow \overline{\mathbb{R}}$ to be convex, proper, and lower semicontinuous on Hilbert spaces X and Y and $K \in \mathcal{L}(X; Y)$ to be a bounded linear operator. We are particularly interested in block-separable

$$(GF) \quad G(x) = \sum_{j=1}^m G_j(P_j x), \quad \text{and} \quad F^*(y) = \sum_{\ell=1}^n F_\ell^*(Q_\ell y),$$

where F^* is the Fenchel conjugate of F . The operators P_1, \dots, P_m are projections in X with $\sum_{j=1}^m P_j = I$ and $P_j P_i = 0$ if $i \neq j$. Likewise, Q_1, \dots, Q_n are projection operators in Y . We assume all the component functions G_j and F_ℓ^* to be convex, proper, and lower semicontinuous, and the subdifferential sum rule to hold for the expressions (GF).

Several first-order optimisation methods have been developed for (P_0) without block-separable structure, typically both G and F convex and K linear. Recently also some non-convexity and

*ModeMat, Escuela Politécnica Nacional, Quito, Ecuador; *previously* Department of Mathematical Sciences, University of Liverpool, United Kingdom. tuomo.valkonen@iki.fi

[†]V3: 2017-11-06, V2: 2017-03-16, V1: 2016-09-16

non-linearity has been introduced [4, 21, 23, 37]. In applications to image processing and data science, one of G or F is typically non-smooth. Effective algorithms operating directly on the primal problem (P_0), or its dual, therefore tend to be a form of classical forward–backward splitting, occasionally called iterative soft-thresholding [1, 12].

In big data optimisation several forward–backward block-coordinate descent methods have been developed for (P_0) with block-separable G . On each step, the methods update only a random subset of blocks $x_j := P_j x$ in parallel; see the review [39] and the original articles [2, 9, 11, 16, 22, 24, 28–31, 42]. Typically F is assumed smooth, and, often, each G_j strongly convex. Besides parallelism, an advantage of these methods is the exploitation of *blockwise* factors of smoothness and strong convexity. These can help convergence by being better than the global factor.

Unfortunately, primal-only and dual-only stochastic methods, as discussed above, are rarely applicable to image processing problems. These, and many other problems, do not satisfy the assumed separability and smoothness assumptions. On the other hand, additional Moreau–Yosida (aka. Huber, aka. Nesterov) regularisation of the problem, which would provide the required smoothness, would alter the problem, losing essential non-smooth characteristics. Generally, even without the splitting of the problem into blocks and the introduction of stochasticity, primal-only or dual-only methods can be inefficient on more complicated problems. Proximal steps, which are typically used to deal with non-smooth components of the problem, can in particular be as expensive as the original optimisation problems itself. In order to make these steps cheap, the problem has to be formulated appropriately. Such a reformulation can often be provided through primal–dual approaches.

With the Fenchel conjugate F^* , we can write (P_0) as

$$(1.1) \quad \min_x \max_y G(x) + \langle Kx, y \rangle - F^*(y).$$

The method of Chambolle and Pock [6, 27] is popular for this formulation. It is also called the PDHGM (Primal-Dual Hybrid Gradient Method, Modified) in [14] and the PDPS (Primal–Dual Proximal Splitting) in [34]. It consists of alternating proximal steps on x and y combined with an over-relaxation step to ensure convergence. The method is closely related to the classical ADMM and Douglas–Rachford splitting. The acronym PDHGM arises from the earlier PDHG [43] that is convergent only in special cases [18]. These connections are discussed in [14].

While early block-coordinate methods only worked with a primal or a dual variable, recently stochastic primal–dual approaches based on the ADMM and the PDHGM have been proposed [3, 15, 25, 26, 33, 40, 41]. Moreover, variants of the ADMM that deterministically update multiple blocks in parallel and afterwards combine the results for the Lagrange multiplier update have been introduced [20]. As with the primal- or dual-only methods, these algorithms can improve convergence by exploiting local properties of the problem. Besides [33, 40, 41] that have restrictive smoothness and strong convexity requirements, little is known about convergence rates.

*In this paper, we will derive block-coordinate descent variants of the PDHGM with known convergence rates: $O(1/N^2)$ if each G_j is strongly convex, $O(1/N)$ without any strong convexity, and mixed $O(1/N^2) + O(1/N)$ if some of the G_j are strongly convex. These rates apply to an ergodic duality gap and strongly convex blocks of the iterates. Our methods have the novelty of blockwise-adapted step lengths. In the imaging applications of Section 5 we will even employ pixelwise-adapted step lengths. Moreover, we can update random subsets of *both* primal and*

dual blocks under a light “nesting condition” on the sampling scheme. Such “doubly-stochastic” updates have previously been possible only in very limited settings [40].

Our present work is based on [37] on the acceleration of the PDHGM when G is strongly convex only on a subspace: the deterministic two-block case $m = 2$ and $n = 1$ of (GF). Besides enabling (doubly-)stochastic updates and an arbitrary number of both primal *and* dual blocks, in the present work, we derive simplified step length rules through a more careful analysis.

The more abstract basis of our present work has been split out in [35]. There we study preconditioning of abstract proximal point methods and “testing” by suitable operators as means of obtaining convergence rates. We recall the relevant aspects of this theory through the course of Sections 2 and 3. In the first of these sections, we start by going through notation and previous research on the PDHGM in more detail. Then we develop the rough structure of our proposed method. This will depend on several structural conditions that we introduce in Section 2. Afterwards in Section 3 we develop convergence estimates based on technical conditions on the various step length and testing parameters. These conditions need to be verified through the development of explicit parameter update rules. We do this in Section 4 along with proving the claimed convergence rates (Theorem 4.5 and its corollaries). We also present there the final, detailed, versions of our proposed algorithms: Algorithm 1 (doubly stochastic) and Algorithm 2 (simplified). We finish with numerical experiments in Section 5.

2 BACKGROUND AND OVERALL STRUCTURE OF THE ALGORITHM

To make the notation definite, we write $\mathcal{L}(X; Y)$ for the space of bounded linear operators between Hilbert spaces X and Y . The identity operator we denote by I . For $T, S \in \mathcal{L}(X; X)$, we use $T \geq S$ to mean that $T - S$ is positive semi-definite; in particular $T \geq 0$ means that T is positive semi-definite. Also for possibly non-self-adjoint T , we introduce the inner product and norm-like notations

$$(2.1) \quad \langle x, z \rangle_T := \langle Tx, z \rangle, \quad \text{and} \quad \|x\|_T := \sqrt{\langle x, x \rangle_T},$$

the latter only defined for positive semi-definite T . We write $T \simeq T'$ if $\langle x, x \rangle_{T'-T} = 0$ for all x .

We denote by $C(X)$ the set of convex, proper, lower semicontinuous functionals from a Hilbert space X to $\overline{\mathbb{R}} := [-\infty, \infty]$. With $G \in C(X)$, $F^* \in C(Y)$, and $K \in \mathcal{L}(X; Y)$, we then wish to solve the minimax problem

$$(P) \quad \min_{x \in X} \max_{y \in Y} G(x) + \langle Kx, y \rangle - F^*(y),$$

assuming the existence of a solution $\hat{u} = (\hat{x}, \hat{y})$ satisfying the optimality conditions

$$(OC) \quad -K^*\hat{y} \in \partial G(\hat{x}), \quad \text{and} \quad K\hat{x} \in \partial F^*(\hat{y}).$$

For the stochastic aspects of our work, we denote by $(\Omega, \mathcal{O}, \mathbb{P})$ the *probability space* consisting of the set Ω of possible realisation of a random experiment, by \mathcal{O} a σ -algebra on Ω , and by \mathbb{P} a probability measure on (Ω, \mathcal{O}) . We denote the expectation corresponding to \mathbb{P} by \mathbb{E} , the conditional probability with respect to a sub- σ -algebra $\mathcal{O}' \subset \mathcal{O}$ by $\mathbb{P}[\cdot | \mathcal{O}']$, and the conditional expectation by $\mathbb{E}[\cdot | \mathcal{O}']$. We refer to [32] for more details.

We also use the next non-standard notation: If \mathcal{O} is a σ -algebra on the space Ω , we denote by $\mathcal{R}(\mathcal{O}; V)$ the space of V -valued random variables A , such that $A : \Omega \rightarrow V$ is \mathcal{O} -measurable.

2.1 PRECONDITIONED PROXIMAL POINT METHODS; TESTING FOR RATES

We use the notation

$$u = (x, y)$$

to combine the primal variable x and dual variable y into a single variable u . Following [19, 37], the primal–dual method of Chambolle and Pock [6] (PDHGM) may then be written in proximal point form as

$$(PP_0) \quad 0 \in H(u^{i+1}) + L_i(u^{i+1} - u^i)$$

for a monotone operator H encoding the optimality conditions (OC) as $0 \in H(\widehat{u})$, and a *preconditioning* or *step length operator* $L_i = L_i^0$. These are

$$(2.2) \quad H(u) := \begin{pmatrix} \partial G(x) + K^*y \\ \partial F^*(y) - Kx \end{pmatrix}, \quad \text{and} \quad L_i^0 := \begin{pmatrix} \tau_i^{-1} & -K^* \\ -\omega_i K & \sigma_{i+1}^{-1} \end{pmatrix}.$$

Here $\tau_i, \sigma_{i+1} > 0$ are step length parameters, and $\omega_i > 0$ an over-relaxation parameter. In the basic version of the algorithm, $\omega_i = 1$, $\tau_i \equiv \tau_0$, and $\sigma_i \equiv \sigma_0$, assuming $\tau_0 \sigma_0 \|K\|^2 < 1$. Observe that we may equivalently parametrise the algorithm by τ_0 and $\delta = 1 - \|K\|^2 \tau_0 \sigma_0 > 0$. The method has $O(1/N)$ rate for the ergodic duality gap that we will return to in Section 3.1.

If G is strongly convex with factor $\gamma > 0$, we may for $\widetilde{\gamma} \in (0, \gamma]$ accelerate

$$(2.3) \quad \omega_i := 1/\sqrt{1 + 2\widetilde{\gamma}\tau_i}, \quad \tau_{i+1} := \tau_i \omega_i, \quad \text{and} \quad \sigma_{i+1} := \sigma_i / \omega_i.$$

This gives $O(1/N^2)$ convergence of $\|x^N - \widehat{x}\|^2$ to zero. If $\widetilde{\gamma} \in (0, \gamma/2]$, we also obtain $O(1/N^2)$ convergence of an ergodic duality gap.

In [37], we extended the PDHGM to partially strongly convex problems: in (GF) this corresponded to the primal two-block and dual single-block case $m = 2$ and $n = 1$ with only G_1 assumed strongly convex. This extension was based on taking in (PP₀) the preconditioner

$$(2.4) \quad L_i = \begin{pmatrix} T_i^{-1} & -K^* \\ -\omega_i K & \Sigma_{i+1}^{-1} \end{pmatrix}$$

for invertible $T_i = \tau_{1,i}P_1 + \tau_{2,i}P_2 \in \mathcal{L}(X; X)$ and $\Sigma_{i+1} = \sigma_{i+1}I \in \mathcal{L}(Y; Y)$. After simple rearrangements of (PP₀), the resulting algorithm could be written more explicitly as

$$(2.5a) \quad x^{i+1} := (I + T_i \partial G)^{-1}(x^i - T_i K^* y^i),$$

$$(2.5b) \quad y^{i+1} := (I + \Sigma_{i+1} \partial F^*)^{-1}(y^i + \Sigma_{i+1} K((1 + \omega_i)x^{i+1} - \omega_i x^i)).$$

Since G is as assumed separable, the first, primal update, splits into separate updates for $x_1^{i+1} := P_1 x^{i+1}$ and $x_2^{i+1} := P_2 x^{i+1}$. Note that this explicit form of the algorithm does not require T_i and Σ_{i+1} to be invertible, unlike (PP₀) with the choice (2.4), so suggests we could develop stochastic methods that randomly choose one, two, or no primal blocks to update.

To study convergence, it is, however, more practical to work with implicit formulations, such as (PP₀). We will shortly see how this works. To make (PP₀) work with non-invertible T_i and Σ_{i+1} , let us reformulate it slightly. In fact, let us define

$$(2.6) \quad W_{i+1} := \begin{pmatrix} T_i & 0 \\ 0 & \Sigma_{i+1} \end{pmatrix}, \quad \text{and (for the moment)} \quad M_{i+1} = \begin{pmatrix} I & -T_i K^* \\ -\widetilde{\omega}_i \Sigma_{i+1} K & I \end{pmatrix}.$$

With this, whether or not T_i and Σ_{i+1} are invertible, (2.5) can be written as the preconditioned proximal point iteration

$$(PP) \quad W_{i+1}H(u^{i+1}) + M_{i+1}(u^{i+1} - u^i) \ni 0,$$

This will be the abstract form of the algorithms that we will develop, however, with the exact form of T_{i+1} , Σ_{i+1} , and M_{i+1} still to be refined.

To study the convergence of (PP), we apply to the *testing* framework introduced in [35, 37]. The idea is to apply $\langle \cdot, u^{i+1} - \widehat{u} \rangle_{Z_{i+1}}$ for a *testing operator* Z_{i+1} to (PP) to “test” it. Thus

$$(2.7) \quad 0 \in \langle W_{i+1}H(u^{i+1}) + M_{i+1}(u^{i+1} - u^i), u^{i+1} - \widehat{u} \rangle_{Z_{i+1}}.$$

We need $Z_{i+1}M_{i+1}$ to be self-adjoint and positive semi-definite. This guarantees that $Z_{i+1}M_{i+1}$ can be used to form the local semi-norm $\| \cdot \|_{Z_{i+1}M_{i+1}}$. Indeed, assuming for some linear operator Ξ_{i+1} that H has the operator-relative (strong) monotonicity property

$$(2.8) \quad \langle H(u') - H(u), u' - u \rangle_{Z_{i+1}W_{i+1}} \geq \|u - u'\|_{Z_{i+1}\Xi_{i+1}}^2 \quad (u, u' \in X \times Y),$$

then a simple application of Pythagoras’ identity

$$\langle u^{i+1} - u^i, u^{i+1} - \widehat{u} \rangle_{Z_{i+1}M_{i+1}} = \frac{1}{2} \|u^{i+1} - u^i\|_{Z_{i+1}M_{i+1}}^2 - \frac{1}{2} \|u^i - \widehat{u}\|_{Z_{i+1}M_{i+1}}^2 + \frac{1}{2} \|u^{i+1} - \widehat{u}\|_{Z_{i+1}M_{i+1}}^2$$

yields

$$\frac{1}{2} \|u^{i+1} - \widehat{u}\|_{Z_{i+1}(M_{i+1} + 2\Xi_{i+1})}^2 + \frac{1}{2} \|u^{i+1} - u^i\|_{Z_{i+1}M_{i+1}}^2 \leq \frac{1}{2} \|u^i - \widehat{u}\|_{Z_{i+1}M_{i+1}}^2.$$

If $Z_{i+2}M_{i+2} \leq Z_{i+1}(M_{i+1} + 2\Xi_{i+1})$ for all i , then summing over $i = 0, \dots, N-1$ gives

$$(2.9) \quad \frac{1}{2} \|u^N - \widehat{u}\|_{Z_{N+1}M_{N+1}}^2 + \sum_{i=0}^{N-1} \frac{1}{2} \|u^{i+1} - u^i\|_{Z_{i+1}M_{i+1}}^2 \leq \frac{1}{2} \|u^0 - \widehat{u}\|_{Z_1M_1}^2.$$

We therefore see that $Z_{i+1}M_{i+1}$ measures the rates of convergence of the iterates. If our iterations are stochastic, to obtain deterministic estimates, we can simply take the expectation in (2.9). However, to obtain estimates on a duality gap, we need to work significantly more. We will, therefore, in the beginning of Section 3, after introducing all the relevant concepts and finalising the setup for the present work, quote the appropriate results from [35].

2.2 STOCHASTIC AND DETERMINISTIC BLOCK UPDATES

We want to update any subset of any number of primal and dual blocks stochastically. Compatible with the separable structure (GF) of G and F^* , we therefore construct from individual (possibly random) step length and testing parameters, $\tau_{j,i}, \sigma_{\ell,i+1} \geq 0$ and $\phi_{j,i}, \psi_{\ell,i+1} > 0$, as well as random subsets $S(i) \subset \{1, \dots, m\}$ and $V(i+1) \subset \{1, \dots, n\}$ the step length and testing operators

$$\begin{aligned}
\text{(S.a)} \quad W_{i+1} &:= \begin{pmatrix} T_i & 0 \\ 0 & \Sigma_{i+1} \end{pmatrix} \quad \text{and} & Z_{i+1} &:= \begin{pmatrix} \Phi_i & 0 \\ 0 & \Psi_{i+1} \end{pmatrix} \quad \text{for} \\
\text{(S.b)} \quad T_i &:= \sum_{j \in S(i)} \tau_{j,i} P_j, & \Sigma_{i+1} &:= \sum_{\ell \in V(i+1)} \sigma_{\ell,i+1} Q_\ell, \\
\text{(S.c)} \quad \Phi_i &:= \sum_{j=1}^m \phi_{j,i} P_j, \quad \text{and} & \Psi_{i+1} &:= \sum_{\ell=1}^n \psi_{\ell,i+1} Q_\ell \quad (i \geq 0).
\end{aligned}$$

We moreover take as the preconditioner

$$\begin{aligned}
\text{(S.d)} \quad M_{i+1} &:= \begin{pmatrix} I & -\Phi_i^{-1} \Lambda_i^* \\ -\Psi_{i+1}^{-1} \Lambda_i & I \end{pmatrix} \quad \text{for} & \Lambda_i &:= K \mathring{T}_i^* \Phi_i^* - \Psi_{i+1} \mathring{\Sigma}_{i+1} K \quad \text{with} \\
\text{(S.e)} \quad \mathring{T}_i &:= \sum_{j \in \mathring{S}(i)} \tau_{j,i} P_j, & \mathring{S}(i) &\subset S(i), \\
\text{(S.f)} \quad \mathring{\Sigma}_{i+1} &:= \sum_{\ell \in \mathring{V}(i+1)} \sigma_{\ell,i+1} Q_\ell, \quad \text{and} & \mathring{V}(i+1) &\subset V(i+1).
\end{aligned}$$

The subsets $S(i)$ and $V(i+1)$ are the indices of the blocks

$$(2.10) \quad x_j := P_j x, \quad \text{and} \quad y_\ell := Q_\ell y$$

of the variables x and y that are to be updated on iteration i .¹ Hence T_i and Σ_{i+1} will not be invertible unless we update all the blocks. Clearly Φ_i , Ψ_{i+1} , T_i , and Σ_{i+1} are self-adjoint and positive semi-definite with Φ_i and Ψ_{i+1} invertible. The subsets $\mathring{S}(i)$ and $\mathring{V}(i+1)$ indicate those blocks of x^{i+1} and of y^{i+1} that are to be updated ‘‘independently’’ of the other variable. We will explain these subsets and the choice of Λ_i in more detail in [Section 2.3](#).

The iterate $u^{i+1} = (x^{i+1}, y^{i+1})$ has to be computable based on random sampling at iteration i and the information gathered (random variable realisations) before commencing the iteration. For the algorithm to be realisable, it cannot depend on the future. We therefore need to be explicit about the space of each random variable. We model the information available just before commencing iteration i by the σ -algebra \mathcal{O}_{i-1} . Thus $\mathcal{O}_{i-1} \subset \mathcal{O}_i$. More precisely, \mathcal{O}_i is the smallest sub- σ -algebra of \mathcal{O} satisfying for all $k = 0, \dots, i$, $j = 1, \dots, m$, and $\ell = 1, \dots, n$ that

$$\begin{aligned}
\text{(R.a)} \quad \tau_{j,k}, \sigma_{\ell,k+1} &\in \mathcal{R}(\mathcal{O}_i; [0, \infty)), & \phi_{j,k}, \psi_{\ell,k+1} &\in \mathcal{R}(\mathcal{O}_i; (0, \infty)), \\
\text{(R.b)} \quad S(k) &\in \mathcal{R}(\mathcal{O}_i; \mathcal{P}(\{1, \dots, m\})), & V(k+1) &\in \mathcal{R}(\mathcal{O}_i; \mathcal{P}(\{1, \dots, n\})), \\
\text{(R.c)} \quad \mathring{S}(k) &\in \mathcal{R}(\mathcal{O}_i; \mathcal{P}(\{1, \dots, m\})), \quad \text{and} & \mathring{V}(k+1) &\in \mathcal{R}(\mathcal{O}_i; \mathcal{P}(\{1, \dots, n\})).
\end{aligned}$$

Here and only here \mathcal{P} denotes the power set. Any other variables can only be random by being constructed from these variables. We thus deduce from [\(S\)](#), and [\(PP\)](#) that

$$\begin{aligned}
T_k &\in \mathcal{R}(\mathcal{O}_i; \mathcal{L}(X; X)), & \Phi_k &\in \mathcal{R}(\mathcal{O}_i; \mathcal{L}(X; X)) & x^{i+1} &\in \mathcal{R}(\mathcal{O}_i; X), \\
\Sigma_{k+1} &\in \mathcal{R}(\mathcal{O}_i; \mathcal{L}(Y; Y)), & \Psi_{k+1} &\in \mathcal{R}(\mathcal{O}_i; \mathcal{L}(Y; Y)), \quad \text{and} & y^{i+1} &\in \mathcal{R}(\mathcal{O}_i; Y).
\end{aligned}$$

¹The iteration index is off-by-one for $\sigma_{\ell,i+1}$ and $\psi_{\ell,i+1}$ for reasons of historical development of the Chambolle–Pock method, when it was not written as a preconditioned proximal point method.

We will also need to assume the *nesting conditions* on sampling,

$$(\mathcal{V}.a) \quad \mathcal{V}(\dot{S}(i)) \cap \dot{V}(i+1) = \emptyset, \quad \mathcal{V}(S(i) \setminus \dot{S}(i)) \cap (V(i+1) \setminus \dot{V}(i+1)) = \emptyset,$$

$$(\mathcal{V}.b) \quad \dot{S}(i) \cup \mathcal{V}^{-1}(\dot{V}(i+1)) \subset S(i), \quad \text{and} \quad \dot{V}(i+1) \cup \mathcal{V}(\dot{S}(i)) \subset V(i+1),$$

where the set

$$(2.11) \quad \mathcal{V}(j) := \{\ell \in \{1, \dots, n\} \mid Q_\ell K P_j \neq 0\}.$$

consists of the dual blocks that are “connected” by K to the primal block with index j . Vice versa, $\mathcal{V}^{-1}(\ell)$ consists of the primal blocks that are “connected” by K to the dual block with index ℓ . Thus $(\mathcal{V}.b)$ states that the independent updates $(\dot{S}(i)$ and $\dot{V}(i+1))$ must propagate from primal to dual and vice versa as non-independent updates $(S(i)$ and $V(i+1))$. The condition $(\mathcal{V}.a)$ restricts connections between primal and dual updates: the first part says that the independently updated blocks cannot be connected. By the second part neither can non-independent updates. If we use $(\mathcal{V}.b)$ as an equality to define $S(i)$ and $V(i+1)$, then the second part of $(\mathcal{V}.a)$ holds if $\mathcal{V}(\mathcal{V}^{-1}(\dot{V}(i+1))) \cap \mathcal{V}(\dot{S}(i)) = \emptyset$, that is, the condition restricts second-degree connections between the independently updated blocks.

To facilitate referring to all the above structural conditions, we introduce:

Assumption 2.1 (main structural condition). We assume the structure (GF) and (S) with the the limitations (\mathcal{R}) and (\mathcal{V}) on randomness.

Clearly

$$(2.12) \quad Z_{i+1} M_{i+1} = \begin{pmatrix} \Phi_i & -\Lambda_i^* \\ -\Lambda_i & \Psi_{i+1} \end{pmatrix}$$

is self-adjoint. We need to prove that it is positive semi-definite. We will do this in [Section 4](#) using the functions κ_ℓ introduced next. As we show afterwards in [Example 2.1](#), these functions are a block structure adapted generalisation of the simple bound $K \leq \|K\|I$.

Definition 2.2. Let $\mathcal{P} := \{P_1, \dots, P_m\}$, and $\mathcal{Q} := \{Q_1, \dots, Q_n\}$. We write $(\kappa_1, \dots, \kappa_n) \in \mathcal{K}(K, \mathcal{P}, \mathcal{Q})$ if each $\kappa_\ell : [0, \infty)^m \rightarrow [0, \infty)$ is monotone ($\ell = 1, \dots, n$) and the following hold:

(i) (Estimation) For all $(z_{\ell,1}, \dots, z_{\ell,m}) \subset [0, \infty)^m$ and $\ell = 1, \dots, n$ the estimate

$$\sum_{j=1}^m \sum_{\ell,k=1}^n z_{\ell,j}^{1/2} z_{k,j}^{1/2} Q_\ell K P_j K^* Q_k \leq \sum_{\ell=1}^n \kappa_\ell(z_{\ell,1}, \dots, z_{\ell,m}) Q_\ell.$$

(ii) (Boundedness) For some $\bar{\kappa} > 0$ and all $(z_1, \dots, z_m) \subset [0, \infty)^m$ the bound

$$\kappa_\ell(z_1, \dots, z_m) \leq \bar{\kappa} \sum_{j=1}^m z_j.$$

(iii) (Non-degeneracy) There exists $\underline{\kappa} > 0$ and for all $j = 1, \dots, m$ a choice of $\ell^*(j) \in \{1, \dots, n\}$ such that for all $(z_1, \dots, z_m) \subset [0, \infty)^m$,

$$\underline{\kappa} z_j \leq \kappa_{\ell^*(j)}(z_1, \dots, z_m) \quad (j = 1, \dots, m).$$

The choice of κ allows us to construct different algorithms. Here we consider a few possibilities, first an easy one, and then a more optimal one.

Example 2.1 (Worst-case κ). We may estimate

$$\sum_{j=1}^m \sum_{\ell, k=1}^n z_{\ell, j}^{1/2} z_{k, j}^{1/2} Q_{\ell} K P_j K^* Q_k \leq \sum_{\ell, k=1}^n \bar{z}_{\ell}^{-1/2} \bar{z}_k^{-1/2} Q_{\ell} K K^* Q_k \leq \sum_{\ell=1}^n \bar{z}_{\ell} \|K\|^2 Q_{\ell}.$$

Therefore [Definition 2.2 \(i\)](#) and [\(ii\)](#) hold with $\bar{\kappa} = \|K\|^2$ for the monotone choice

$$\kappa_{\ell}(z_1, \dots, z_m) := \|K\|^2 \max\{z_1, \dots, z_m\}.$$

Clearly also $\underline{\kappa} = \bar{\kappa}$ for any choice of $\ell^*(j) \in \{1, \dots, n\}$. This choice of κ_{ℓ} corresponds to the rule $\tau \sigma \|K\|^2 < 1$ in the PDHGM method.

Example 2.2 (Balanced κ). Take minimal κ_{ℓ} satisfying [Definition 2.2](#) and the balancing condition $\kappa_{\ell}(z_{\ell, 1}, \dots, z_{\ell, m}) = \kappa_k(z_{k, 1}, \dots, z_{k, m})$, $(\ell, k = 1, \dots, n)$. This requires problem-specific analysis, but tends to perform well, as we will see in [Section 5](#).

2.3 JUSTIFICATION OF THE PRECONDITIONER AND SAMPLING RESTRICTIONS

With M_{i+1} of the form [\(S.d\)](#) for any Λ_i , the implicit method [\(PP\)](#) expands as

$$(2.13a) \quad 0 \in T_i \partial G(x^{i+1}) + T_i K^* y^{i+1} + (x^{i+1} - x^i) - \Phi_i^{-1} \Lambda_i^* (y^{i+1} - y^i), \quad \text{and}$$

$$(2.13b) \quad 0 \in \Sigma_{i+1} \partial F^*(y^{i+1}) - \Sigma_{i+1} K x^{i+1} - \Psi_{i+1}^{-1} \Lambda_i (x^{i+1} - x^i) + (y^{i+1} - y^i).$$

This can easily be rearranged as

$$(2.14a) \quad x^{i+1} := (I + T_i \partial G)^{-1} (x^i + \Phi_i^{-1} \Lambda_i^* (y^{i+1} - y^i) - T_i K^* y^{i+1}),$$

$$(2.14b) \quad y^{i+1} := (I + \Sigma_{i+1} \partial F^*)^{-1} (y^i + \Psi_{i+1}^{-1} \Lambda_i (x^{i+1} - x^i) + \Sigma_{i+1} K x^{i+1}).$$

This method is still not explicit as the primal and dual updates potentially depend on each other. We will now show how the removal of this dependency leads to our choice of Λ_i in [\(S.d\)](#).

Indeed, due to the compatible block-separable structures [\(S\)](#) and [\(GF\)](#), multiplying [\(2.13a\)](#) by P_j for $j = 1, \dots, m$, and [\(2.13b\)](#) by Q_{ℓ} for $\ell = 1, \dots, n$, [\(2.14\)](#) can be split blockwise as

$$x_j^{i+1} = (I + \chi_{S(i)}(j) \tau_{j, i} \partial G_j)^{-1} (x_j^i + P_j [\Phi_i^{-1} \Lambda_i^* (y^{i+1} - y^i) - T_i K^* y^{i+1}]) \quad (j = 1, \dots, m),$$

$$y_{\ell}^{i+1} = (I + \chi_{V(i+1)}(\ell) \sigma_{\ell, i+1} \partial F_{\ell}^*)^{-1} (y_{\ell}^i + Q_{\ell} [\Psi_{i+1}^{-1} \Lambda_i (x^{i+1} - x^i) + \Sigma_{i+1} K x^{i+1}]) \quad (\ell = 1, \dots, n).$$

For $S(i)$ and $V(i+1)$ to have the intended meaning that only the corresponding blocks are updated, we need to choose Λ_i such that

$$(2.15) \quad x_j^{i+1} = x_j^i \quad (j \notin S(i)), \quad \text{and} \quad y_{\ell}^{i+1} = y_{\ell}^i \quad (\ell \notin V(i+1)).$$

Since $P_j T_i = 0$ for $j \notin S(i)$, and $Q_\ell \Sigma_{i+1} = 0$ for $\ell \notin V(i+1)$, this is to say

$$(2.16a) \quad P_j \Phi_i^{-1} \Lambda_i^* (y^{i+1} - y^i) = 0 \quad (j \notin S(i)), \quad \text{and}$$

$$(2.16b) \quad Q_\ell \Psi_{i+1}^{-1} \Lambda_i (x^{i+1} - x^i) = 0 \quad (\ell \notin V(i+1)).$$

Taking $\Lambda_i = K T_i^* \Phi_i^*$ would allow computing x^{i+1} before y^{i+1} and satisfy (2.16a). If we further had $K T_i^* \Phi_i^* = \omega_i \Psi_{i+1} \Sigma_{i+1} K$, then also (2.16b) would hold and (S.d) would reproduce M_{i+1} of (2.6). Symmetrically, $\Lambda_i = -\Sigma_{i+1} \Psi_{i+1} K$ would make y^{i+1} independent of x^{i+1} . Such conditions will, however, rarely be satisfiable unless, deterministically, $S(i) \equiv \{1, \dots, m\}$ and $V(i+1) \equiv \{1, \dots, n\}$. Nevertheless, motivated by this, we designate subsets of blocks of x^{i+1} and y^{i+1} to be updated independently of the other variable. This are the subsets $\mathring{S}(i)$ and $\mathring{V}(i+1)$ in (S.e) and (S.f). Then picking Λ_i as in (S.d) achieves our objective:

Lemma 2.1. *Suppose Assumption 2.1 holds. Then (2.15) and (2.16) hold.*

Proof. We already know that (2.16) implies (2.15). Using (S.d), (2.16) can be rewritten

$$P_j \Phi_i^{-1} [\Phi_i \mathring{T}_i K^* - K^* \mathring{\Sigma}_{i+1}^* \Psi_{i+1}^*] (y^{i+1} - y^i) = 0 \quad (j \notin S(i)), \quad \text{and}$$

$$Q_\ell \Psi_{i+1}^{-1} [K T_i^* \Phi_i^* - \Psi_{i+1} \mathring{\Sigma}_{i+1} K] (x^{i+1} - x^i) = 0 \quad (\ell \notin V(i+1)).$$

Clearly $P_j \mathring{T}_i K^*$ for $j \notin S(i)$. Therefore, the first condition holds if $P_j K^* \mathring{\Sigma}_{i+1}^* = 0$ for $j \notin S(i)$. This is to say $j \notin \mathcal{V}^{-1}(\mathring{V}(i+1))$, which is guaranteed by (V.b). Likewise, the second condition holds if $Q_\ell K T_i^* = 0$ for $\ell \notin V(i+1)$, which is also guaranteed by (V.b). \square

2.4 OVERALL STRUCTURE OF THE PROPOSED METHOD

Setting $T_i^\perp := T_i - \mathring{T}_i$, and $\Sigma_{i+1}^\perp := \Sigma_{i+1} - \mathring{\Sigma}_{i+1}$, we can now rewrite (2.14) as

$$(2.17a) \quad q^{i+1} := \Phi_i^{-1} K^* \mathring{\Sigma}_{i+1}^* \Psi_{i+1}^* (y^{i+1} - y^i) + T_i^\perp K^* y^{i+1},$$

$$(2.17b) \quad x^{i+1} := (I + T_i \partial G)^{-1} (x^i - \mathring{T}_i K^* y^i - q^{i+1}),$$

$$(2.17c) \quad z^{i+1} := \Psi_{i+1}^{-1} K T_i^* \Phi_i^* (x^{i+1} - x^i) + \Sigma_{i+1}^\perp K x^{i+1},$$

$$(2.17d) \quad y^{i+1} := (I + \Sigma_{i+1} \partial F^*)^{-1} (y^i + \mathring{\Sigma}_{i+1} K x^i + z^{i+1}).$$

Due to the first part of (V.a), $P_j q^{i+1} = 0$ and $Q_\ell z^{i+1} = 0$ for $j \in \mathring{S}(i)$ and $\ell \in \mathring{V}(i+1)$. The second part of (V.a) moreover implies $T_i^\perp K^* Q_\ell = 0$ and $\Sigma_{i+1}^\perp K P_j$ for $\ell \in V(i+1) \setminus \mathring{V}(i+1)$ and $j \in S(i) \setminus \mathring{S}(i)$. The first part of (V.a) and (V.b) moreover imply $\Sigma_{i+1}^\perp \Theta_{i+1} = \Sigma_{i+1} \Theta_{i+1} = \Psi_{i+1}^{-1} K T_i^* \Phi_i^*$ and $T_i^\perp B_{i+1}^* = T_i B_{i+1}^* = \Phi_i^{-1} K^* \mathring{\Sigma}_{i+1}^* \Psi_{i+1}^*$ for

$$\Theta_i := \sum_{j \in \mathring{S}(i)} \sum_{\ell \in \mathcal{V}(j)} \theta_{\ell,j,i} Q_\ell K P_j \quad \text{with} \quad \theta_{\ell,j,i+1} := \frac{\tau_{j,i} \phi_{j,i}}{\sigma_{\ell,i+1} \psi_{\ell,i+1}} \quad \text{and}$$

$$B_i := \sum_{\ell \in \mathring{V}(i+1)} \sum_{j \in \mathcal{V}^{-1}(\ell)} b_{\ell,j,i} Q_\ell K P_j \quad \text{with} \quad b_{\ell,j,i+1} := \frac{\sigma_{\ell,i+1} \psi_{\ell,i+1}}{\tau_{j,i} \phi_{j,i}}$$

Setting $\hat{x}^{i+1} := \sum_{j \in \hat{S}(i)} P_j x^{i+1}$ and $\hat{y}^{i+1} := \sum_{\ell \in \hat{V}(i+1)} Q_\ell x^{i+1}$ we therefore obtain

$$(2.18a) \quad q^{i+1} := \Phi_i^{-1} K^* \hat{\Sigma}_{i+1}^* \Psi_{i+1}^* (\hat{y}^{i+1} - y^i) + T_i^\perp K^* \hat{y}^{i+1} = T_i^\perp [B_{i+1}^* (\hat{y}^{i+1} - y^i) + \hat{y}^{i+1}] \quad \text{and}$$

$$(2.18b) \quad z^{i+1} := \Psi_{i+1}^{-1} K \hat{T}_i^* \Phi_i^* (\hat{x}^{i+1} - x^i) + \Sigma_{i+1}^\perp K \hat{x}^{i+1} = \Sigma_{i+1}^\perp [\Theta_{i+1} (\hat{x}^{i+1} - x^i) + \hat{x}^{i+1}].$$

Introducing w^{i+1} and v^{i+1} such that $\Psi_{i+1}^\perp w^{i+1} = z^{i+1}$ and $\Phi_i^\perp v^{i+1} = q^{i+1}$, and using (GF), we can write the method given by (2.17) and (2.18) as

$$(2.19a) \quad \hat{x}^{i+1} := (I + \hat{T}_i \partial G)^{-1} (x^i - \hat{T}_i K^* y^i),$$

$$(2.19b) \quad \hat{y}^{i+1} := (I + \hat{\Sigma}_{i+1} \partial F^*)^{-1} (y^i + \hat{\Sigma}_{i+1} K x^i),$$

$$(2.19c) \quad w^{i+1} := \Theta_{i+1} (\hat{x}^{i+1} - x^i) + \hat{x}^{i+1},$$

$$(2.19d) \quad v^{i+1} := B_{i+1}^* (\hat{y}^{i+1} - y^i) + \hat{y}^{i+1},$$

$$(2.19e) \quad x^{i+1} := (I + T_i^\perp \partial G)^{-1} (\hat{x}^{i+1} - T_i^\perp v^{i+1}),$$

$$(2.19f) \quad y^{i+1} := (I + \Sigma_{i+1}^\perp \partial F^*)^{-1} (\hat{y}^{i+1} + \Sigma_{i+1}^\perp w^{i+1}).$$

Due to (GF), these operations can further be split into blockwise operations with no dependencies on so far uncomputed blocks. We delay doing this explicitly until we are ready to present our final Algorithms 1 and 2 towards the end of the theoretical part of the paper.

We conclude the present structural development by explicitly stating what we have proved in the preceding paragraphs:

Lemma 2.2. *Suppose Assumption 2.1 holds. Then (2.19) is equivalent to (PP).*

3 A BASIC CONVERGENCE ESTIMATE

Now that we have the overall structure of the proposed algorithms established in (2.19), we need to develop rules for the step length and testing parameters that yield a convergent method. This will require, in particular, the positive semi-definiteness of $Z_{i+1} M_{i+1}$, as we recall from the discussion leading up to (2.9). Indeed, since in general, without any strong convexity, we can only obtain gap (and weak) convergence, we need to refine that argument.

In Sections 3.1 to 3.5, we will derive some quite technical conditions that the step length parameters, testing parameters, and block selection probabilities need to satisfy. From these basic estimates, we then develop explicit convergence rates in the next section. In the final Section 3.6, we will also discuss permissible sampling patterns.

3.1 A BOUND ON ERGODIC DUALITY GAPS

Recall the basis of the testing technique, (2.7). In the single-block case ($T_i = \tau_i I$, $\Sigma_{i+1} = \sigma_{i+1} I$, $\Phi_{i+1} = \phi_{i+1} I$, and $\Psi_{i+1} = \psi_{i+1} I$), instead of using $\hat{u} \in H^{-1}(0)$ and the operator-relative monotonicity (2.8) to eliminate H , using the convexity of G and F^* we can also estimate

$$(3.1) \quad \begin{aligned} \langle H(u^{i+1}), u^{i+1} - \hat{u} \rangle_{Z_{i+1} W_{i+1}} &\geq \phi_i \tau_i [G(x^{i+1}) - G(\hat{x})] + \psi_{i+1} \sigma_{i+1} [F^*(y^{i+1}) - F^*(\hat{y})] \\ &\quad + \phi_i \tau_i \langle K^* y^{i+1}, x^{i+1} - \hat{x} \rangle - \psi_{i+1} \sigma_i \langle K x^{i+1}, y^{i+1} - \hat{y} \rangle =: \tilde{\mathcal{G}}_{i+1}. \end{aligned}$$

With this, (2.9) can be improved to

$$\frac{1}{2}\|u^N - \widehat{u}\|_{Z_{N+1}M_{N+1}}^2 + \sum_{i=0}^{N-1} \left(\widetilde{\mathcal{G}}_{i+1} + \frac{1}{2}\|u^{i+1} - u^i\|_{Z_{i+1}M_{i+1}}^2 \right) \leq \frac{1}{2}\|u^0 - \widehat{u}\|_{Z_1M_1}^2.$$

We would like to develop the ‘‘preliminary gaps’’ $\widetilde{\mathcal{G}}_{i+1}$ into a (Lagrangian) duality gap

$$(3.2) \quad \mathcal{G}(x, y) := (G(x) + \langle \widehat{y}, Kx \rangle - F^*(\widehat{y})) - (G(\widehat{x}) + \langle y, K\widehat{x} \rangle - F^*(y)).$$

The first obstacle we face are the differing factors in front of G and F^* . This suggests to impose $\phi_i \tau_i = \psi_{i+1} \sigma_{i+1}$. For the PDHGM, it however turns out that $\phi_i \tau_i = \psi_i \sigma_i$. After taking care of some technical details, this can be dealt with by an index realignment argument [35].

With multiple blocks, we can get a similar estimate as (3.1) with factor the $\phi_{j,i} \tau_{j,i}$ in front of G_j and $\psi_{\ell,i+1} \sigma_{\ell,i+1}$ in front of F_ℓ^* . To derive a gap estimate, the preceding discussion suggests to impose $T_i \Phi_i = \bar{\eta}_i I$ and $\Sigma_{i+1} \Psi_{i+1} = \bar{\eta}_i I$ or $\Sigma_i \Psi_i = \bar{\eta}_i I$ for some scalar $\bar{\eta}_i > 0$. This kind of *coupling* between the blocks will be one of the main restrictions that we face in the development our method. In the stochastic setting, it turns out [35] that we can relax the coupling slightly: do it in expectation. Correspondingly, we assume for some $\bar{\eta}_i > 0$ either

$$(3.3a) \quad \mathbb{E}[T_i^* \Phi_i^*] = \bar{\eta}_i I, \quad \text{and} \quad \mathbb{E}[\Psi_{i+1} \Sigma_{i+1}] = \bar{\eta}_i I, \quad (i \geq 1), \quad \text{or}$$

$$(3.3b) \quad \mathbb{E}[T_i^* \Phi_i^*] = \bar{\eta}_i I, \quad \text{and} \quad \mathbb{E}[\Psi_i \Sigma_i] = \bar{\eta}_i I, \quad (i \geq 1).$$

The second condition is an extension of what we saw the standard PDHGM to satisfy. The first condition, which is off-by-one compared to the second, will, however, be the only alternative that doubly-stochastic methods can satisfy.

A further difficulty with developing (3.1) into a gap estimate are the remaining terms involving K . Even after rearrangements we can only get an ergodic estimate [35]. To express such estimates, corresponding to the conditions (3.3a) and (3.3b), we introduce

$$(3.4) \quad \zeta_N := \sum_{i=0}^{N-1} \bar{\eta}_i \quad \text{and} \quad \zeta_{*,N} := \sum_{i=1}^{N-1} \bar{\eta}_i,$$

and the ergodic sequences

$$(3.5) \quad \widetilde{x}_N := \zeta_N^{-1} \mathbb{E} \left[\sum_{i=0}^{N-1} T_i^* \Phi_i^* x^{i+1} \right], \quad \widetilde{y}_N := \zeta_N^{-1} \mathbb{E} \left[\sum_{i=0}^{N-1} \Sigma_{i+1}^* \Psi_{i+1}^* y^{i+1} \right],$$

$$(3.6) \quad \widetilde{x}_{*,N} := \zeta_{*,N}^{-1} \mathbb{E} \left[\sum_{i=1}^{N-1} T_i^* \Phi_i^* x^{i+1} \right], \quad \widetilde{y}_{*,N} := \zeta_{*,N}^{-1} \mathbb{E} \left[\sum_{i=1}^{N-1} \Sigma_i^* \Psi_i^* y^i \right].$$

The coupling conditions (3.3a) and (3.3b) then produce two different ergodic gaps, $\mathcal{G}(\widetilde{x}_N, \widetilde{y}_N)$ and $\mathcal{G}(\widetilde{x}_{*,N}, \widetilde{y}_{*,N})$. We demonstrate this in the next theorem from [35]. It forms the basis for our work in the remaining sections. The fundamental arguments for the proof are those that led to (2.9), however, the gap estimate requires significant additional technical work.

Theorem 3.1. *Suppose Assumption 2.1 (main structural condition) holds with $Z_{i+1}M_{i+1}$ positive semi-definite. Write $\Gamma := \sum_{j=1}^m \gamma_j P_j$ for $\gamma_j \geq 0$ the factor of (strong) convexity of G_j . With $\tilde{\Gamma} = \sum_{j=1}^m \tilde{\gamma}_j P_j \in \mathcal{L}(X; X)$, assuming one of the following alternatives to hold, let*

$$(3.7) \quad \tilde{g}_N := \begin{cases} 0, & 0 \leq \tilde{\Gamma} \leq \Gamma, \\ \zeta_N \mathcal{G}(\tilde{x}_N, \tilde{y}_N), & 0 \leq \tilde{\Gamma} \leq \Gamma/2; \text{ (3.3a) holds,} \\ \zeta_{*,N} \mathcal{G}(\tilde{x}_{*,N}, \tilde{y}_{*,N}), & 0 \leq \tilde{\Gamma} \leq \Gamma/2; \text{ (3.3b) holds.} \end{cases}$$

Also define

$$\Xi_{i+1}(\tilde{\Gamma}) := \begin{pmatrix} 2T_i \tilde{\Gamma} & 2T_i K^* \\ -2\Sigma_{i+1} K & 0 \end{pmatrix} \quad \text{and} \quad D_{i+1}(\tilde{\Gamma}) := Z_{i+2} M_{i+2} - Z_{i+1} (\Xi_{i+1}(\tilde{\Gamma}) + M_{i+1}).$$

Then the iterates $u^i = (x^i, y^i)$ of (PP) satisfy for any $\hat{u} \in H^{-1}(0)$ the estimate

$$(3.8) \quad \frac{1}{2} \mathbb{E} [\|u^N - \hat{u}\|_{Z_N M_N}^2] + \tilde{g}_N \leq \frac{1}{2} \|u^0 - \hat{u}\|_{Z_1 M_1}^2 + \sum_{i=0}^{N-1} \frac{1}{2} \mathbb{E} [\|u^{i+1} - \hat{u}\|_{D_{i+1}(\tilde{\Gamma})}^2 - \|u^{i+1} - u^i\|_{Z_{i+1} M_{i+1}}^2].$$

Proof. This is [35, Theorem 5.5] with $\Delta_{i+1}(\tilde{\Gamma}) := \frac{1}{2} \|u^{i+1} - \hat{u}\|_{D_{i+1}(\tilde{\Gamma})}^2 - \frac{1}{2} \|u^{i+1} - u^i\|_{Z_{i+1} M_{i+1}}^2$ and the condition $\tilde{\Gamma} = \Gamma$ relaxed to $0 \leq \tilde{\Gamma} \leq \Gamma$, which is possible because if g_j is strongly convex with factor $\gamma_j > 0$, it is strongly convex with any smaller non-negative factor. Moreover, [35, Example 5.1] shows that the blockwise structure (GF), (S) has an ergodic convexity property that produces the gaps $\mathcal{G}(\tilde{x}_N, \tilde{y}_N)$ and $\mathcal{G}(\tilde{x}_{*,N}, \tilde{y}_{*,N})$ \square

In the standard PDHGM we can ensure $D_{i+1}(\tilde{\Gamma}) \simeq 0$ [35]. However, in our present setting, we will not generally be able to enforce this, so these operators will introduce a penalty in (3.8). A lot of our remaining work will consist of controlling this penalty. We also need to estimate from below and show that $Z_N M_N$ is positive semi-definite.

3.2 NOTATIONS AND ASSUMPTIONS

For convenience, we introduce

$$\begin{aligned} \hat{\tau}_{j,i} &:= \tau_{j,i} \chi_{S(i)}(j), & \hat{\sigma}_{\ell,i} &:= \sigma_{\ell,i} \chi_{V(i)}(\ell), \\ \pi_{j,i} &:= \mathbb{P}[j \in S(i) \mid \mathcal{O}_{i-1}], & \nu_{\ell,i+1} &:= \mathbb{P}[\ell \in V(i+1) \mid \mathcal{O}_{i-1}], \\ \dot{\pi}_{j,i} &:= \mathbb{P}[j \in \dot{S}(i) \mid \mathcal{O}_{i-1}], & \dot{\nu}_{\ell,i+1} &:= \mathbb{P}[\ell \in \dot{V}(i+1) \mid \mathcal{O}_{i-1}]. \end{aligned}$$

The first two denote ‘‘effective’’ step lengths on iteration i , while the rest are shorthands for the probabilities of the primal block j or the dual block ℓ being contained in the corresponding set on iteration i . Recalling (S.d), we also write

$$(3.9) \quad \Lambda_i = \sum_{j=1}^m \sum_{\ell \in \mathcal{V}(j)} \lambda_{\ell,j,i} Q_\ell K P_j \quad \text{with} \quad \lambda_{\ell,j,i} := \phi_{j,i} \hat{\tau}_{j,i} \chi_{\dot{S}(i)}(j) - \psi_{\ell,i+1} \hat{\sigma}_{\ell,i+1} \chi_{\dot{V}(i+1)}(\ell).$$

We require the following technical assumption, which we will verify through explicit step length and testing parameter update rule development in the next section. We indicate the rough intended use of each condition in parentheses after the statement.

Assumption 3.1 (step length parameter restrictions). We assume for each $i \in \mathbb{N}$ the following, constants independent of i , and the *same alternatives* holding for each i :

- (a) We are given $(\kappa_1, \dots, \kappa_n) \in \mathcal{K}(K, \mathcal{P}, \mathcal{Q})$ (see [Definition 2.2](#)), and for some $\delta \in (0, 1)$,

$$(1 - \delta)\psi_{\ell, i+1} \geq \kappa_{\ell}(\lambda_{\ell, 1, i}^2 \phi_{1, i}^{-1}, \dots, \lambda_{\ell, m, i}^2 \phi_{m, i}^{-1}) \quad (\ell = 1, \dots, n).$$

(This condition generalises the condition $\tau\sigma\|K\|^2 < 1$ for the standard PDHGM, needed to ensure the positivity of the local metric $Z_{i+1}M_{i+1}$.)

- (b) We are given $\eta_i \in \mathcal{R}(\mathcal{O}_{i-1}; (0, \infty))$ and $\eta_{\tau, i}^{\perp}, \eta_{\sigma, i}^{\perp} \in \mathcal{R}(\mathcal{O}_{i-1}; [0, \infty))$ such that $\eta_{i+1} \geq \eta_i$,

$$\eta_i \cdot \min_j (\pi_{j, i} - \hat{\pi}_{j, i}) \geq \eta_{\tau, i}^{\perp}, \quad \text{and} \quad \eta_{i+1} \cdot \min_{\ell} (v_{\ell, i+1} - \hat{v}_{\ell, i+1}) \geq \eta_{\sigma, i}^{\perp}.$$

(This is needed to annihilate the off-diagonal of D_{i+1} in the penalty term.)

- (c) Either

(c-i) $\mathbb{E}[\eta_{\tau, i}^{\perp} - \eta_{\sigma, i}^{\perp}] = \text{constant}$; or

(c-ii) $\eta_{\tau, i}^{\perp} = 0$ and $\eta_{\sigma, i}^{\perp} = \eta_{i+1}$.

(These are needed to ensure the coupling conditions [\(3.3a\)](#) or [\(3.3b\)](#), respectively.)

- (d) The step lengths parameters

$$(3.10a) \quad \tau_{j, i} = \begin{cases} \frac{\eta_i - \phi_{j, i-1} \tau_{j, i-1} \chi_{S(i-1) \setminus \hat{S}(i-1)}(j)}{\phi_{j, i} \hat{\pi}_{j, i}}, & j \in \hat{S}(i), \\ \frac{\eta_{\tau, i}^{\perp}}{\phi_{j, i} (\pi_{j, i} - \hat{\pi}_{j, i})}, & j \in S(i) \setminus \hat{S}(i), \end{cases}$$

$$(3.10b) \quad \sigma_{j, i+1} = \begin{cases} \frac{\eta_i - \psi_{j, i} \sigma_{j, i} \chi_{V(i) \setminus \hat{V}(i)}(j)}{\psi_{j, i+1} \hat{v}_{\ell, i+1}}, & j \in \hat{V}(i+1), \\ \frac{\eta_{\sigma, i}^{\perp}}{\psi_{j, i+1} (v_{\ell, i+1} - \hat{v}_{\ell, i+1})}, & j \in V(i+1) \setminus \hat{V}(i+1). \end{cases}$$

For $i = 0$ we take $\tau_{j, -1} := 0$ and $\sigma_{j, 0} := 0$.

(This rule is also needed to annihilate the off-diagonal of D_{i+1} in the penalty term.)

- (e) Let $\gamma_j \geq 0$ by the factor of (strong) convexity of G_j , and $\tilde{\gamma}_j \in [0, \gamma_j]$, ($j = 1, \dots, m$). Also let $\alpha_i > 0$ and define

$$(3.11a) \quad q_{j, i+2}(\tilde{\gamma}_j) := (\mathbb{E}[\phi_{j, i+1} - \phi_{j, i}(1 + 2\hat{\tau}_{j, i}\tilde{\gamma}_j) | \mathcal{O}_i] + \alpha_i |\mathbb{E}[\phi_{j, i+1} - \phi_{j, i}(1 + 2\hat{\tau}_{j, i}\tilde{\gamma}_j) | \mathcal{O}_i]| - \delta \phi_{j, i}) \chi_{S(i)}(j), \quad \text{and}$$

$$(3.11b) \quad h_{j, i+2}(\tilde{\gamma}_j) := \mathbb{E}[\phi_{j, i+1} - \phi_{j, i}(1 + 2\hat{\tau}_{j, i}\tilde{\gamma}_j) | \mathcal{O}_{i-1}] + \alpha_i^{-1} |\mathbb{E}[\phi_{j, i+1} - \phi_{j, i}(1 + 2\hat{\tau}_{j, i}\tilde{\gamma}_j) | \mathcal{O}_i]|.$$

Then for some $C_x > 0$ either

$$(3.12a) \quad \|x_j^{i+1} - \widehat{x}_j\|^2 \leq C_x \quad (j = 1, \dots, m) \quad \text{or}$$

$$(3.12b) \quad h_{j,i+2}(\widetilde{Y}_j) \leq 0 \quad \text{and} \quad q_{j,i+2}(\widetilde{Y}_j) \leq 0 \quad (j = 1, \dots, m),$$

(This is needed to bound the primal components in the penalty term.)

(f) For some $C_y > 0$ either

$$(3.13a) \quad \mathbb{E}[\psi_{\ell,i+2} - \psi_{\ell,i+1} | \mathcal{O}_i] \geq 0, \quad \|y_\ell^{i+1} - \widehat{y}_\ell\|^2 \leq C_y \quad (\ell = 1, \dots, n) \quad \text{or}$$

$$(3.13b) \quad \mathbb{E}[\psi_{\ell,i+2} - \psi_{\ell,i+1} | \mathcal{O}_i] = 0 \quad (\ell = 1, \dots, n).$$

(This is needed to bound the dual components in the penalty term.)

It is important that [Assumption 3.1](#) is consistent with [Assumption 2.1](#), in particular that the step lengths generated by the former are non-negative. We will prove this in [Lemma 3.4](#). Before this, we start the main goal of the present section, the following specialisation of [Theorem 3.1](#).

Proposition 3.2. *Suppose [Assumption 2.1](#) (main structural condition) and [Assumption 3.1](#) (step length restrictions) hold. Then the iterates of (PP) satisfy for any $\widehat{u} \in H^{-1}(0)$ the estimate*

$$(3.14) \quad \sum_{j=1}^m \frac{\delta}{2\mathbb{E}[\phi_{j,N}^{-1}]} \cdot \mathbb{E}[\|x_j^N - \widehat{x}_j\|^2] + \widetilde{g}_N \leq \frac{1}{2} \|u^0 - \widehat{u}\|_{Z_0 M_0}^2 + \sum_{j=1}^m \frac{1}{2} d_{j,N}^x(\widetilde{Y}_j) + \sum_{\ell=1}^n \frac{1}{2} d_{\ell,N}^y,$$

where

$$(3.15a) \quad \widetilde{g}_N := \begin{cases} \zeta_N \mathcal{G}(\widetilde{x}_N, \widetilde{y}_N), & \text{case (c-i) and } \widetilde{y}_j \leq \gamma_j/2 \text{ for all } j, \\ \zeta_{*,N} \mathcal{G}(\widetilde{x}_{*,N}, \widetilde{y}_{*,N}), & \text{case (c-ii) and } \widetilde{y}_j \leq \gamma_j/2 \text{ for all } j, \\ 0, & \text{otherwise,} \end{cases}$$

$$(3.15b) \quad d_{j,N}^x(\widetilde{Y}_j) := \sum_{i=0}^{N-1} \delta_{j,i+2}^x(\widetilde{Y}_j) \quad d_{\ell,N}^y := \sum_{i=0}^{N-1} \delta_{\ell,i+2}^y,$$

$$(3.15c) \quad \delta_{j,i+2}^x(\widetilde{Y}_j) := 4C_x \mathbb{E}[\max\{0, q_{j,i+2}(\widetilde{Y}_j)\}] + C_x \mathbb{E}[\max\{0, h_{j,i+2}(\widetilde{Y}_j)\}], \quad \text{and}$$

$$(3.15d) \quad \delta_{\ell,i+2}^y := 9C_y \mathbb{E}[\psi_{\ell,i+2} - \psi_{\ell,i+1}].$$

Proof. We use [Lemma 3.7](#) or [Lemma 3.8](#) (to follow) to verify one of the coupling conditions (3.3a) or (3.3b). Then we obtain (3.8) from [Theorem 3.1](#). Next, we use [Lemmas 3.3](#) and [3.6](#) (to follow) to estimate $Z_{N+1} M_{N+1} \geq \begin{pmatrix} \delta \Phi_N & 0 \\ 0 & 0 \end{pmatrix}$ and

$$\mathbb{E}[\|u^{i+1} - \widehat{u}\|_{D_{i+1}(\widetilde{\Gamma})}^2 - \|u^{i+1} - u^i\|_{Z_{i+1} M_{i+1}}^2] \leq \sum_{j=1}^m \delta_{j,i+2}^x(\widetilde{Y}_j) + \sum_{\ell=1}^n \delta_{\ell,i+2}^y.$$

Therefore (3.8) yields

$$\frac{\delta}{2} \mathbb{E}[\|x^N - \widehat{x}\|_{\Phi_N}^2] + \widetilde{g}_N \leq \frac{1}{2} \|u^0 - \widehat{u}\|_{Z_0 M_0}^2 + \frac{1}{2} \sum_{i=0}^{N-1} \left(\sum_{j=1}^m \delta_{j,i+2}^x(\widetilde{Y}_j) + \sum_{\ell=1}^n \delta_{\ell,i+2}^y \right).$$

By Hölder's inequality

$$\mathbb{E}[\|x^N - \widehat{x}\|_{\Phi_N}^2] = \sum_{j=1}^m \mathbb{E}[\phi_{j,N} \|x_j^N - \widehat{x}_j\|^2] \geq \sum_{j=1}^m \mathbb{E}[\|x_j^N - \widehat{x}_j\|^2] / \mathbb{E}[\phi_{j,N}^{-1}].$$

The estimate (3.14) is now immediate. \square

3.3 LOWER BOUND ON THE LOCAL METRIC

Lemma 3.3. *Suppose Assumption 2.1 (main structural condition) and Assumption 3.1(a) hold. Then $Z_{i+1}M_{i+1} \geq \begin{pmatrix} \delta\Phi_i & 0 \\ 0 & 0 \end{pmatrix}$.*

Proof. Since Φ_{i+1} is self-adjoint and positive definite, using (2.12) and Cauchy's inequality, for any $\delta \in (0, 1)$ we deduce

$$(3.16) \quad Z_{i+1}M_{i+1} = \begin{pmatrix} \Phi_i & -\Lambda_i^* \\ -\Lambda_i & \Psi_{i+1} \end{pmatrix} \geq \begin{pmatrix} \delta\Phi_i & 0 \\ 0 & \Psi_{i+1} - \frac{1}{1-\delta}\Lambda_i\Phi_i^{-1}\Lambda_i^* \end{pmatrix}.$$

We therefore require $(1-\delta)\Psi_{i+1} \geq \Lambda_i\Phi_i^{-1}\Lambda_i^*$, which can be expanded as

$$(3.17) \quad (1-\delta) \sum_{\ell=1}^n \psi_{\ell, i+1} Q_\ell \geq \sum_{j=1}^m \sum_{\ell, k=1}^n \lambda_{\ell, j, i} \lambda_{k, j, i} \phi_{j, i}^{-1} Q_\ell K P_j K^* Q_k.$$

This follows from Definition 2.2 (i) with $z_{\ell, j} := \lambda_{\ell, j, i}^2 \phi_{j, i}^{-1}$. \square

3.4 BOUNDS ON THE PENALTY TERMS

The structural setup (S) gives

$$(3.18) \quad D_{i+1}(\widetilde{\Gamma}) = \begin{pmatrix} \Phi_{i+1} - \Phi_i(I + 2T_i\widetilde{\Gamma}) & \Lambda_i^* - \Lambda_{i+1}^* - 2\Phi_i T_i K^* \\ 2\Psi_{i+1}\Sigma_{i+1}K + \Lambda_i - \Lambda_{i+1} & \Psi_{i+2} - \Psi_{i+1} \end{pmatrix} \\ \simeq \begin{pmatrix} \Phi_{i+1} - \Phi_i(I + 2T_i\widetilde{\Gamma}) & A_{i+2}^* \\ A_{i+2} & \Psi_{i+2} - \Psi_{i+1} \end{pmatrix} \quad \text{for} \\ A_{i+2} := (\Psi_{i+1}\Sigma_{i+1}K - \Lambda_{i+1}) + (\Lambda_i - K T_i^* \Phi_i^*).$$

Lemma 3.4. *Suppose Assumption 2.1 (main structural condition) and Assumption 3.1(b) & (d) hold. Then*

$$(3.19) \quad \mathbb{E}[A_{i+2}|O_i](x^{i+1} - x^i) = 0, \quad \mathbb{E}[A_{i+2}^*|O_i](y^{i+1} - y^i) = 0, \quad \mathbb{E}[A_{i+2}^*|O_{i-1}] = 0.$$

Moreover, if $\phi_{j, i}, \psi_{\ell, i+1} > 0$ for all $i \in \mathbb{N}$, then $\tau_{j, i}, \sigma_{\ell, i+1} \geq 0$ for all $i \in \mathbb{N}$. In particular, Assumption 3.1 is consistent with Assumption 2.1 requiring $\tau_{j, i}, \sigma_{\ell, i+1} \geq 0$ and $\phi_{j, i}, \psi_{\ell, i+1} > 0$ for all $i \in \mathbb{N}$; $j = 1, \dots, m$; and $\ell = 1, \dots, n$.

Proof. We start by claiming that

$$(3.20) \quad \mathbb{E}[\lambda_{\ell,j,i+1}|\mathcal{O}_i] = \psi_{\ell,i+1}\hat{\sigma}_{\ell,i+1}(1 - \chi_{\dot{V}(i+1)}(\ell)) - \phi_{j,i}\hat{\tau}_{j,i}(1 - \chi_{\dot{S}(i)}(j))$$

whenever $\ell \in \mathcal{V}(j)$. Indeed, inserting (3.9) into (3.20), we see the former to be satisfied if (for any given η_{i+1}),

$$(3.21a) \quad \mathbb{E}[\phi_{j,i+1}\hat{\tau}_{j,i+1}\chi_{\dot{S}(i+1)}(j)|\mathcal{O}_i] = \eta_{i+1} - \phi_{j,i}\hat{\tau}_{j,i}(1 - \chi_{\dot{S}(i)}(j)) \geq 0, \quad \text{and}$$

$$(3.21b) \quad \mathbb{E}[\psi_{\ell,i+2}\hat{\sigma}_{\ell,i+2}\chi_{\dot{V}(i+2)}(\ell)|\mathcal{O}_i] = \eta_{i+1} - \psi_{\ell,i+1}\hat{\sigma}_{\ell,i+1}(1 - \chi_{\dot{V}(i+1)}(\ell)) \geq 0,$$

over $j = 1, \dots, m$; $\ell = 1, \dots, n$; and $i \geq -1$, taking $\dot{S}(-1) = \{1, \dots, m\}$ and $\dot{V}(0) = \{1, \dots, n\}$.

We can also write (3.21a) as

$$(3.22) \quad \mathbb{E}[\phi_{j,i+1}\hat{\tau}_{j,i+1}\chi_{\dot{S}(i+1)}(j)|\mathcal{O}_i] = \eta_{i+1} - \phi_{j,i}\tau_{j,i}\chi_{S(i)\setminus\dot{S}(i)}(j) \geq 0.$$

If $j \notin S(i) \setminus \dot{S}(i)$, since $\eta_{i+1} \geq 0$ by Assumption 3.1(b), it is clear that the inequality in (3.22) holds. If $j \in S(i) \setminus \dot{S}(i)$, using the corresponding case of (3.10a), we rewrite the inequality as $\eta_{i+1} \geq \eta_{\tau,i}^{\perp}/(\pi_{j,i} - \pi_{j,i}^{\circ})$. This is verified by Assumption 3.1(b). Comparing to Assumption 3.1(d), the inequality in (3.22) now inductively verifies, as claimed, $\tau_{j,i+1} \geq 0$ for all $i \in \mathbb{N}$ provided $\phi_{j,i} > 0$ for all $i \in \mathbb{N}$.

To verify the equality in (3.22), let $\mathcal{O}_i^+ \supset \mathcal{O}_i$ be the smallest σ -algebra also containing the set $\{\omega \in \Omega \mid j \in \dot{S}(\omega)(i+1)\}$ (now not abusing notation for random variables, with ω standing for the random realisation that we typically omit). By Assumption 3.1(d), more precisely (3.10b) shifted from i to $i+1$, we see that $\phi_{j,i+1}\tau_{j,i+1}$ is \mathcal{O}_i^+ -measurable. Therefore, by standard properties of conditional expectations (see, e.g., [32])

$$(3.23) \quad \begin{aligned} \mathbb{E}[\phi_{j,i+1}\hat{\tau}_{j,i+1}\chi_{\dot{S}(i+1)}(j)|\mathcal{O}_i] &= \mathbb{E}[\mathbb{E}[\phi_{j,i+1}\hat{\tau}_{j,i+1}\chi_{\dot{S}(i+1)}(j)|\mathcal{O}_i^+]| \mathcal{O}_i] \\ &= \mathbb{E}[\mathbb{E}[\phi_{j,i+1}\tau_{j,i+1}|\mathcal{O}_i^+]| \mathcal{O}_i] \\ &= \mathbb{E}[\mathbb{E}[1|\mathcal{O}_i^+]\phi_{j,i+1}\tau_{j,i+1}|\mathcal{O}_i] = \mathbb{E}[\pi_{j,i+1}^{\circ}\phi_{j,i+1}\tau_{j,i+1}|\mathcal{O}_i]. \end{aligned}$$

Further expanding with (3.10b) shifted from i to $i+1$, and d using $\eta_{i+1} \in \mathcal{R}(\mathcal{O}_i; (0, \infty))$ from Assumption 3.1(b), we obtain

$$(3.24) \quad \mathbb{E}[\pi_{j,i+1}^{\circ}\phi_{j,i+1}\tau_{j,i+1}|\mathcal{O}_i] = \mathbb{E}[\eta_{i+1} - \phi_{j,i}\tau_{j,i}\chi_{S(i)\setminus\dot{S}(i)}(j)|\mathcal{O}_i] = \eta_{i+1} - \phi_{j,i}\tau_{j,i}\chi_{S(i)\setminus\dot{S}(i)}(j).$$

This verifies the equality in (3.22). Thus (3.21a) holds.

Similarly we can verify (3.21b) and $\sigma_{\ell,i+1} \geq 0$. Thus (3.20) holds, as does the non-negativity claim on the dual step lengths.

Using (\mathcal{V} .b) and (3.20), we now observe that $\lambda_{\ell,j,i}$ satisfies

$$(3.25a) \quad \lambda_{\ell,j,i} = 0, \quad (j \notin S(i) \text{ or } \ell \notin V(i+1)), \quad \text{and}$$

$$(3.25b) \quad \mathbb{E}[\lambda_{\ell,j,i+1}|\mathcal{O}_i] = \tilde{\lambda}_{\ell,j,i+1}, \quad (j = 1, \dots, m; \ell \in \mathcal{V}(j)),$$

for $\tilde{\lambda}_{\ell,j,i+1} := \psi_{\ell,i+1}\hat{\sigma}_{\ell,i+1} + \lambda_{\ell,j,i} - \phi_{j,i}\hat{\tau}_{j,i}$. Using (2.15), which follows from Lemma 2.1, (3.19) expands as

$$(3.26a) \quad \mathbb{E}[\lambda_{\ell,j,i+1}|\mathcal{O}_i] = \tilde{\lambda}_{\ell,j,i+1} \quad (j \in S(i), \ell \in \mathcal{V}(j)),$$

$$(3.26b) \quad \mathbb{E}[\lambda_{\ell,j,i+1}|\mathcal{O}_i] = \tilde{\lambda}_{\ell,j,i+1} \quad (\ell \in V(i+1), j \in \mathcal{V}^{-1}(\ell)), \quad \text{and}$$

$$(3.26c) \quad \mathbb{E}[\lambda_{\ell,j,i+1}|\mathcal{O}_{i-1}] = \mathbb{E}[\tilde{\lambda}_{\ell,j,i+1}|\mathcal{O}_{i-1}], \quad (j = 1, \dots, m; \ell \in \mathcal{V}(j)).$$

Clearly (3.25b) implies (3.26a) and (3.26b). Moreover, applying $\mathbb{E}[\cdot|\mathcal{O}_{i-1}]$ to (3.25b) and using standard properties of nested conditional expectations we obtain (3.26c). We have therefore verified (3.19). \square

Corollary 3.5. *Suppose Assumption 3.1 (b) & (d) hold. Then*

$$\begin{aligned} \mathbb{E}[\phi_{j,i+1}\hat{\tau}_{j,i+1}|\mathcal{O}_i] &= \eta_{i+1} + \eta_{\tau,i+1}^\perp - \eta_{\tau,i}^\perp, \quad \text{and} \\ \mathbb{E}[\psi_{\ell,i+2}\hat{\sigma}_{\ell,i+2}|\mathcal{O}_i] &= \eta_{i+1} + \eta_{\sigma,i+1}^\perp - \eta_{\sigma,i}^\perp. \end{aligned}$$

Proof. Arguing analogously to (3.23) and (3.24) with the cases $j \in S(i) \setminus \dot{S}(i)$ and $\ell \in V(i+1) \setminus \dot{V}(i+1)$ of Assumption 3.1 (d), we deduce

$$\begin{aligned} \mathbb{E}[\phi_{j,i+1}\hat{\tau}_{j,i+1}(1 - \chi_{\dot{S}(i+1)}(j))|\mathcal{O}_i] &= \eta_{\tau,i+1}^\perp, \quad \text{and} \\ \mathbb{E}[\psi_{\ell,i+2}\hat{\sigma}_{\ell,i+2}(1 - \chi_{\dot{V}(i+2)}(\ell))|\mathcal{O}_i] &= \eta_{\sigma,i+1}^\perp. \end{aligned}$$

Combined with (3.21) (in the proof of Lemma 3.4) these imply the claim. \square

For the next lemma we recall the coordinate notation x_j and y_ℓ from (2.10).

Lemma 3.6. *Suppose Assumption 2.1 (main structural condition) and Assumption 3.1 (step length parameter restrictions) hold. Then*

$$\mathbb{E}[\|u^{i+1} - \hat{u}\|_{D_{i+1}(\tilde{\Gamma})}^2 - \|u^{i+1} - u^i\|_{Z_{i+1}M_{i+1}}^2] \leq \sum_{j=1}^m \delta_{j,i+2}^x(\tilde{Y}_j) + \sum_{\ell=1}^n \delta_{\ell,i+2}^y,$$

where $\delta_{j,i+2}^x(\tilde{Y}_j)$ and $\delta_{\ell,i+2}^y$ are given in (3.15c) and (3.15d), respectively.

Proof. Since $u^{i+1} \in \mathcal{R}(\mathcal{O}_i; X \times Y)$ and $u^i \in \mathcal{R}(\mathcal{O}_{i-1}; X \times Y)$, standard nesting properties of conditional expectations show

$$(3.27) \quad \begin{aligned} \mathbb{E}[\|u^{i+1} - \hat{u}\|_{D_{i+1}(\tilde{\Gamma})}^2] &= \mathbb{E}[\|u^{i+1} - u^i\|_{\mathbb{E}[D_{i+1}(\tilde{\Gamma})|\mathcal{O}_i]}^2 + \|u^i - \hat{u}\|_{\mathbb{E}[D_{i+1}(\tilde{\Gamma})|\mathcal{O}_{i-1}]}^2 \\ &\quad + 2\langle u^{i+1} - u^i, u^i - \hat{u} \rangle_{\mathbb{E}[D_{i+1}(\tilde{\Gamma})|\mathcal{O}_i]}]. \end{aligned}$$

By Lemma 3.4, (3.19) holds. Using (3.18), we therefore expand (3.27) into

$$\begin{aligned} \mathbb{E}[\|u^{i+1} - \hat{u}\|_{D_{i+1}(\tilde{\Gamma})}^2] &= \mathbb{E}[\|x^{i+1} - x^i\|_{\mathbb{E}[\Phi_{i+1}-\Phi_i(I+2T_i\tilde{\Gamma})|\mathcal{O}_i]}^2 + \|x^i - \hat{x}\|_{\mathbb{E}[\Phi_{i+1}-\Phi_i(I+2T_i\tilde{\Gamma})|\mathcal{O}_{i-1}]}^2 \\ &\quad + \|y^{i+1} - y^i\|_{\mathbb{E}[\Psi_{i+2}-\Psi_{i+1}|\mathcal{O}_i]}^2 + \|y^i - \hat{y}\|_{\mathbb{E}[\Psi_{i+2}-\Psi_{i+1}|\mathcal{O}_{i-1}]}^2 \\ &\quad + 2\langle x^{i+1} - x^i, x^i - \hat{x} \rangle_{\mathbb{E}[\Phi_{i+1}-\Phi_i(I+2T_i\tilde{\Gamma})|\mathcal{O}_i]} + 2\langle y^{i+1} - y^i, y^i - \hat{y} \rangle_{\mathbb{E}[\Psi_{i+2}-\Psi_{i+1}|\mathcal{O}_i]}]. \end{aligned}$$

By [Assumption 3.1 \(f\)](#), $\mathbb{E}[\Psi_{i+2} - \Psi_{i+1} | \mathcal{O}_i] \geq 0$. Standard properties of conditional expectations guarantee $\mathbb{E}[\mathbb{E}[\Psi_{i+2} - \Psi_{i+1} | \mathcal{O}_i] | \mathcal{O}_{i-1}] = \mathbb{E}[\Psi_{i+2} - \Psi_{i+1} | \mathcal{O}_{i-1}]$. By [Lemma 3.3](#), moreover

$$-\|u^{i+1} - u^i\|_{Z_{i+1}M_{i+1}}^2 \leq -\delta \|x^{i+1} - x^i\|_{\Phi_i}.$$

Use of Cauchy's inequality for arbitrary factors $\alpha_i, \beta_i > 0$ therefore yields

$$\begin{aligned} & \mathbb{E} \left[\|u^{i+1} - \widehat{u}\|_{D_{i+1}(\widetilde{\Gamma})}^2 - \|u^{i+1} - u^i\|_{Z_{i+1}M_{i+1}}^2 \right] \\ &= \mathbb{E} \left[\|x^{i+1} - x^i\|_{\mathbb{E}[\Phi_{i+1} - \Phi_i(I+2T_i\widetilde{\Gamma}) | \mathcal{O}_i] + \alpha_i \mathbb{E}[\Phi_{i+1} - \Phi_i(I+2T_i\widetilde{\Gamma}) | \mathcal{O}_i]}^2 \right. \\ & \quad + \|x^i - \widehat{x}\|_{\mathbb{E}[\Phi_{i+1} - \Phi_i(I+2T_i\widetilde{\Gamma}) | \mathcal{O}_{i-1}] + \alpha_i^{-1} \mathbb{E}[\Phi_{i+1} - \Phi_i(I+2T_i\widetilde{\Gamma}) | \mathcal{O}_i]}^2 \\ & \quad \left. + (1 + \beta_i) \|y^{i+1} - y^i\|_{\mathbb{E}[\Psi_{i+2} - \Psi_{i+1} | \mathcal{O}_i]}^2 + (1 + \beta_i^{-1}) \|y^i - \widehat{y}\|_{\mathbb{E}[\Psi_{i+2} - \Psi_{i+1} | \mathcal{O}_{i-1}]}^2 \right]. \end{aligned}$$

Here we write $|\sum_{j=1}^m c_j P_j| := \sum_{j=1}^m |c_j| P_j$. Therefore, choosing $\beta_i = 1/2$, splitting the estimates into blocks, and using [Assumption 3.1 \(e\) & \(f\)](#), we obtain the claim. \square

It is relatively easy to satisfy [Assumption 3.1 \(f\)](#) and to bound $\delta_{\ell, i+2}^y$. To estimate $\delta_{j, i+2}^x(\widetilde{Y}_j)$, we need to derive more involved update rules. We next construct one example.

Example 3.1 (Random primal test updates). If (3.12a) holds, take $\rho_j \geq 0$, otherwise take $\rho_j = 0$ ($j = 1, \dots, m$). Set

$$(3.28) \quad \phi_{j, i+1} := \phi_{j, i} (1 + 2\widetilde{Y}_j \hat{\tau}_{j, i}) + 2\rho_j \pi_{j, i}^{-1} \chi_{S(i)}(j), \quad (j = 1, \dots, m; i \in \mathbb{N}).$$

Then it is not difficult to show that $\phi_{j, i+1} \in \mathcal{R}(\mathcal{O}_i; (0, \infty))$ and $\delta_{j, i+2}^x(\widetilde{Y}_j) = 18C_x \rho_j$.

If we set $\rho_j = 0$ and have just a single deterministically updated block, (3.28) is the standard rule (2.3) with $\phi_i = \tau_i^{-2}$. The role of $\rho_j > 0$ is to ensure some (slower) acceleration on non-strongly-convex blocks with $\widetilde{Y}_j = 0$. This is necessary for convergence rate estimates.

The difficulty with (3.28) is that the coupling parameter η_{i+1} will depend on the random realisations of $S(i)$ through $\phi_{j, i+1}$. This will require communication in a parallel implementation of the algorithm. We therefore desire to update $\phi_{j, i+1}$ deterministically. We delay the introduction of an appropriate update rule to [Section 4](#).

3.5 SATISFACTION OF THE COUPLING CONDITIONS

We still need to satisfy either of the coupling conditions (3.3) to obtain gap estimates.

Lemma 3.7. *Suppose [Assumption 2.1](#) (main structural condition), [Assumption 3.1 \(d\)](#), (b) & (c-i) hold. Then the coupling condition (3.3a) holds.*

Proof. The condition (3.3a) holds if $\mathbb{E}[\phi_{j, i+1} \hat{\tau}_{j, i+1}] = \bar{\eta}_{i+1} = \mathbb{E}[\psi_{\ell, i+2} \hat{\sigma}_{\ell, i+2}]$ for some $\bar{\eta}_{i+1}$ for all $j = 1, \dots, m$ and $\ell = 1, \dots, n$. Taking $\bar{\eta}_{i+1} := \mathbb{E}[\eta_{i+1} + \eta_{\tau, i+1}^\perp - \eta_{\tau, i}^\perp]$, the claim follows from [Corollary 3.5](#) and [Assumption 3.1 \(c-i\)](#). \square

The alternative coupling condition (3.3b) requires $\mathbb{E}[\phi_{j,i+1}\hat{\tau}_{j,i+1}] = \bar{\eta}_{i+1} = \mathbb{E}[\psi_{\ell,i+1}\hat{\sigma}_{\ell,i+1}]$ for some $\bar{\eta}_{i+1}$. By Corollary 3.5, this holds when

$$(3.29) \quad \mathbb{E}[\eta_{i+1} + \eta_{\tau,i+1}^\perp - \eta_{\tau,i}^\perp] = \bar{\eta}_i = \mathbb{E}[\eta_i + \eta_{\sigma,i}^\perp - \eta_{\sigma,i-1}^\perp].$$

It is not clear how to satisfy this simultaneously with Assumption 3.1 (c-i), so we use (c-ii).

Lemma 3.8. *Suppose Assumption 2.1 (main structural condition), Assumption 3.1 (d) & (c-ii) hold. Then (b) holds if and only if $\mathring{V}(i+1) = \emptyset$ and $V(i+1) = \{1, \dots, n\}$. When this is the case, the coupling condition (3.3b) holds, necessarily $S(i) = \mathring{S}(i)$, and*

$$(3.30a) \quad \tau_{j,i} = \eta_i / (\phi_{j,i} \pi_{j,i}^\circ) \quad (j \in S(i)),$$

$$(3.30b) \quad \sigma_{j,i+1} = \eta_{i+1} / \psi_{j,i+1} \quad (j \in \mathcal{V}(S(i))).$$

Proof. Assumption 3.1 (c-ii), i.e., $\eta_{\tau,i}^\perp = 0$ and $\eta_{\sigma,i}^\perp = \eta_{i+1}$ reduces (b) to $\min_\ell (v_{\ell,i+1} - \mathring{v}_{\ell,i+1}) \geq 1$. This holds if and only if $v_{\ell,i+1} \equiv 1$ and $\mathring{v}_{\ell,i+1} \equiv 0$ for all $\ell = 1, \dots, m$. This holds, as claimed, if and only if $\mathring{V}(i+1) = \emptyset$, $V(i+1) = \{1, \dots, n\}$. Clearly in this case (\mathcal{V}) holds if and only if $S(i) = \mathring{S}(i)$ this implies (\mathcal{V}). With Assumption 3.1 (b) verified, Corollary 3.5 now rewrites (3.3b) as (3.29). This is clearly verified by $\eta_{\tau,i}^\perp = 0$ and $\eta_{\sigma,i}^\perp = \eta_{i+1}$. Finally, (3.30) is a specialisation of Assumption 3.1 (d) to the choices of (c-ii). \square

Remark 3.9. *We had to impose full dual updates to satisfy (3.3b). This is akin to most existing primal–dual coordinate descent methods [3, 15, 33]. The algorithms in [25, 26, 40] are more closely related to our method, however, only [40] provides convergence rates for single-block sampling schemes under full strong convexity of both G and F^* .*

3.6 SAMPLING PATTERNS

There are not many possible fully deterministic sampling patterns allowed by (\mathcal{V}) with Assumption 3.1. Indeed, (3.21a) reads in the deterministic setting

$$\phi_{j,i+1}\tau_{j,i+1}\chi_{\mathring{S}(i+1)}(j) + \phi_{j,i}\hat{\tau}_{j,i}\chi_{S(i)\setminus\mathring{S}(i)}(j) = \eta_{i+1}.$$

Since $\eta_{i+1} > 0$, $j \notin S(i) \setminus \mathring{S}(i)$ implies $j \in \mathring{S}(i+1)$, which implies $j \notin S(i+i) \setminus \mathring{S}(i+1)$. Therefore, once in the independently updated set, the block j will always stay there. Due to (\mathcal{V} .b), if $\mathring{V}(i+1) \neq \emptyset$ consistently, for most K , the set $S(i)$ will grow. Therefore, after a small number of iterations N , either $j \in \mathring{S}(i)$ for $i \geq N$, or $j \in S(i) = \{1, \dots, n\}$. Similar considerations hold for the dual blocks. Therefore, the way each block is updated in deterministic methods is, after a small number of iterations, fixed. There does not, therefore, appear to be significant improvements possible over consistently taking $S(i) = \mathring{S}(i) = \{1, \dots, m\}$, $\mathring{V}(i+1) = \emptyset$ and $V(i+1) = \{1, \dots, m\}$ (or the converse dual-first order).

Regarding stochastic algorithms, we start with a few options for sampling $S(i)$ in Algorithm 2 with iteration-independent probabilities $\pi_{j,i} \equiv \pi_j$.

Example 3.2 (Independent probabilities). If all the blocks $\{1, \dots, m\}$ are chosen independently, we have $\mathbb{P}(\{j, k\} \subset S(i)) = \pi_j \pi_k$ for $j \neq k$, where $\pi_j \in (0, 1]$.

Example 3.3 (Fixed number of random blocks). If we have a fixed number M of processors, we may want to choose a subset $S(i) \subset \{1, \dots, m\}$ such that $\#S(i) = M$.

The next example gives a simple way to satisfy $(\mathcal{V}.a)$ for [Algorithm 1](#).

Example 3.4 (Alternating x-y and y-x steps). Let us randomly alternate between $\mathring{S}(i) = \emptyset$ and $\mathring{V}(i+1) = \emptyset$. That is, with some probability \mathbb{p}_x , we choose to take an x-y step that omits lines 9 and 8 in [Algorithm 1](#), and with probability $1 - \mathbb{p}_x$, an y-x step that omits the lines 7 and 10. If $\tilde{\pi}_j = \mathbb{P}[j \in \mathring{S} | \mathring{S} \neq \emptyset]$, and $\tilde{v}_\ell = \mathbb{P}[\ell \in \mathring{V} | \mathring{V} \neq \emptyset]$ denote the probabilities of the rule used to sample $\mathring{S} = \mathring{S}(i)$ and $\mathring{V} = \mathring{V}(i+1)$ when non-empty, then (\mathcal{V}) gives

$$\begin{aligned} \mathring{\pi}_j &= \mathbb{p}_x \tilde{\pi}_j, & \pi_j &= \mathbb{p}_x \tilde{\pi}_j + (1 - \mathbb{p}_x) \mathbb{P}[j \in \mathcal{V}^{-1}(\mathring{V}) | \mathring{V} \neq \emptyset], \\ \mathring{v}_\ell &= (1 - \mathbb{p}_x) \tilde{v}_\ell, & v_\ell &= (1 - \mathbb{p}_x) \tilde{v}_\ell + \mathbb{p}_x \mathbb{P}[\ell \in \mathcal{V}(\mathring{S}) | \mathring{S} \neq \emptyset]. \end{aligned}$$

To compute π_j and v_ℓ we thus need to know \mathcal{V} and the exact sampling pattern.

Remark 3.10. Based on [Example 3.4](#), we can derive an algorithm where the only randomness comes from alternating between full x-y and full y-x steps.

4 RATES OF CONVERGENCE

We now need to satisfy [Assumption 3.1](#). This involves choosing update rules for η_{i+1} , $\eta_{\tau, i+1}^\perp$, $\eta_{\sigma, i+1}^\perp$, $\phi_{j, i+1}$ and $\psi_{\ell, i+1}$. At the same time, to obtain good convergence rates, we need to make $d_{j, N}^x(\tilde{y}_j)$ and $d_{\ell, N}^y$ small in [\(3.14\)](#). We do these tasks here, including stating two final versions of our algorithm [Algorithm 1](#) (doubly stochastic) and [Algorithm 2](#) (full dual updates). Specifically, in [Section 4.1](#) we introduce and study a deterministic alternative to the example random update rule for $\phi_{j, i+1}$ in [Example 3.1](#). The analysis of the new rule is easier, and it allows the computation of η_i , which will also be deterministic, without communication in parallel implementations of our algorithms. Afterwards, in [Section 4.2](#) we look at possible choices for the parameters $\eta_{\tau, i}^\perp$ and $\eta_{\sigma, i}^\perp$, which are only needed in stochastic variants of [Algorithm 1](#). In [Sections 4.3](#) to [4.6](#) we then give various useful choices of η_i and $\psi_{\ell, i}$ that yield concrete convergence rates.

We assume for simplicity that the sampling pattern is independent of iteration,

$$(4.1) \quad \mathring{\pi}_{j, i} \equiv \mathring{\pi}_j > 0, \quad \mathring{v}_{\ell, i} \equiv \mathring{v}_\ell, \quad \pi_{j, i} \equiv \pi_j, \quad \text{and} \quad v_{\ell, i} \equiv v_\ell.$$

4.1 DETERMINISTIC PRIMAL TEST UPDATES

The next lemma gives a deterministic alternative to [Example 3.1](#). We recall that $\gamma_j \geq 0$ is the factor of (strong) convexity of G_j .

Lemma 4.1. *Suppose Assumption 3.1 (b) & (d), and (4.1) hold, and that $i \mapsto \eta_{\tau,i}^\perp$ is non-decreasing. Suppose, moreover, that either (3.12a) holds or $\sup_{j=1,\dots,m} \rho_j = 0$. Also take $\tau_{j,0}, \phi_{j,0} > 0$ and $\bar{\gamma}_j \geq 0$ such that $\rho_j + \bar{\gamma}_j > 0$, and set*

$$(4.2) \quad \phi_{j,i+1} := \phi_{j,i} + 2(\bar{\gamma}_j \eta_i + \rho_j), \quad (j = 1, \dots, m; i \in \mathbb{N}).$$

Then for some $c_j > 0$ and all $N \geq 1$ holds

$$(4.3a) \quad \phi_{j,N+1} \in \mathcal{R}(\mathcal{O}_{N-1}; (0, \infty)),$$

$$(4.3b) \quad \mathbb{E}[\phi_{j,N}] = \phi_{j,0} + 2\rho_j N + 2\bar{\gamma}_j \sum_{i=0}^{N-1} \mathbb{E}[\eta_i], \quad \text{and}$$

$$(4.3c) \quad \mathbb{E}[\phi_{j,N}^{-1}] \leq c_j N^{-1}, \quad (N \geq 1).$$

If $\bar{\gamma}_j \in [\bar{\gamma}_j, \gamma_j]$, ($j = 1, \dots, m$), satisfy

$$(4.3d) \quad 2\bar{\gamma}_j \bar{\gamma}_j \eta_i \leq (\bar{\gamma}_j - \bar{\gamma}_j) \delta \phi_{j,i}, \quad (j \in S(i), i \in \mathbb{N}),$$

then Assumption 3.1 (e) holds, and

$$(4.3e) \quad d_{j,N}^x(\bar{\gamma}_j) = 18\rho_j C_x N.$$

Finally, if $\eta_i \geq b_j \min_j \phi_{j,i}^p$ for some $p, b_j > 0$, then for some $\tilde{c}_j > 0$ holds

$$(4.3f) \quad 1 \geq \bar{\gamma}_j \tilde{c}_j N^{p+1} \mathbb{E}[\phi_{j,N}^{-1}], \quad (N \geq 4).$$

Proof. Since Assumption 3.1 (b) guarantees $\eta_i \in \mathcal{R}(\mathcal{O}_{i-1}; (0, \infty))$, we deduce (4.3a) from (4.2). In fact, $\phi_{j,i+1}$ is deterministic as long as η_i is deterministic.

The claim (4.3b) is immediate from using (4.2) to compute

$$(4.4) \quad \phi_{j,N} = \phi_{j,N-1} + 2(\bar{\gamma}_j \eta_{N-1} + \rho_j) = \phi_{j,0} + 2\rho_j N + 2\bar{\gamma}_j \sum_{i=0}^{N-1} \eta_i.$$

Since $i \mapsto \eta_i$ is non-decreasing, clearly $\phi_{j,N} \geq 2N\bar{\rho}_j$ for $\bar{\rho}_j := \rho_j + \bar{\gamma}_j \eta_0 > 0$. Then $\phi_{j,N}^{-1} \leq \frac{1}{2\bar{\rho}_j N}$. Taking the expectation proves (4.3c).

Clearly (4.3f) holds if $\bar{\gamma}_j = 0$, so assume $\bar{\gamma}_j > 0$. Using the assumption $\eta_i \geq b_j \min_j \phi_{j,i}^p$ and $\phi_{j,i} \geq 2i\bar{\rho}_j$ that we just proved in (4.4), we estimate

$$\phi_{j,N} \geq \phi_{j,0} + b_j (2\bar{\rho}_j)^p \sum_{i=1}^N i^p \geq \phi_{j,0} + b_j (2\bar{\rho}_j)^p \int_2^N x^p dx \geq \phi_{j,0} + p^{-1} b_j (2\bar{\rho}_j)^p (N^{p+1} - 2).$$

Thus $\phi_{j,N}^{-1} \leq 1/(\bar{\gamma}_j \tilde{c}_j N^{1+p})$ for some $\tilde{c}_j > 0$. Taking the expectation proves (4.3f).

It remains to prove (4.3e) and Assumption 3.1 (e). Abbreviating $\gamma_{j,i} := \bar{\gamma}_j + \rho_j \eta_i^{-1}$, we write $\phi_{j,i+1} = \phi_{j,i} + 2\gamma_{j,i} \eta_i$. Since $i \mapsto \eta_{\tau,i}^\perp$ is non-decreasing, Corollary 3.5 gives

$$(4.5) \quad \mathbb{E}[\phi_{j,i} \hat{\tau}_{j,i} | \mathcal{O}_{i-1}] = \eta_i + \eta_{\tau,i}^\perp - \eta_{i-1,\tau}^\perp \geq \eta_i.$$

Expanding the defining equation (3.11b) of $h_{j,i+2}(\bar{\gamma}_j)$, with the help of (4.5) we estimate

$$\begin{aligned} h_{j,i+2}(\bar{\gamma}_j) &= 2\mathbb{E}[\gamma_{j,i}\eta_i - \bar{\gamma}_j\phi_{j,i}\hat{\tau}_{j,i}|\mathcal{O}_{i-1}] + 2\alpha_i^{-1}|\mathbb{E}[\gamma_{j,i}\eta_i - \bar{\gamma}_j\phi_{j,i}\hat{\tau}_{j,i}|\mathcal{O}_i]| \\ &\leq 2(\gamma_{j,i} - \bar{\gamma}_j)\eta_i + 2\alpha_i^{-1}|\gamma_{j,i}\eta_i - \bar{\gamma}_j\phi_{j,i}\hat{\tau}_{j,i}| \\ &\leq 2(1 + \alpha_i^{-1})\rho_j + 2(\bar{\gamma}_j - \bar{\gamma}_j)\eta_i + 2\alpha_i^{-1}|\bar{\gamma}_j\eta_i - \bar{\gamma}_j\phi_{j,i}\hat{\tau}_{j,i}|. \end{aligned}$$

Since (4.3d) implies $\bar{\gamma}_j \leq \bar{\gamma}_j$, if also

$$(4.6) \quad \alpha_i^{-1}|\bar{\gamma}_j\eta_i - \bar{\gamma}_j\phi_{j,i}\hat{\tau}_{j,i}| \leq (\bar{\gamma}_j - \bar{\gamma}_j)\eta_i,$$

then

$$(4.7) \quad \mathbb{E}[\max\{0, h_{j,i+2}(\bar{\gamma}_j)\}] \leq 2(1 + \alpha_i^{-1})\rho_j.$$

We claim (4.6) this to hold for

$$(4.8) \quad \alpha_i := \begin{cases} \min_j \bar{\gamma}_j / (\bar{\gamma}_j - \bar{\gamma}_j), & \bar{\gamma}_j\eta_i > \bar{\gamma}_j\phi_{j,i}\hat{\tau}_{j,i}, \\ \min_j (\bar{\gamma}_j\hat{\pi}_j^{-1} + \bar{\gamma}_j) / (\bar{\gamma}_j - \bar{\gamma}_j), & \bar{\gamma}_j\eta_i \leq \bar{\gamma}_j\phi_{j,i}\hat{\tau}_{j,i}. \end{cases}$$

The case $\bar{\gamma}_j\eta_i > \bar{\gamma}_j\phi_{j,i}\hat{\tau}_{j,i}$ is clear. Otherwise, to justify the case $\bar{\gamma}_j\eta_i \leq \bar{\gamma}_j\phi_{j,i}\hat{\tau}_{j,i}$, we observe that (4.6) can in this case be rewritten as $\bar{\gamma}_j\phi_{j,i}\hat{\tau}_{j,i} \leq (\alpha_i(\bar{\gamma}_j - \bar{\gamma}_j) - \bar{\gamma}_j)\eta_i$. With the choice of α_i in (4.8), we see this to hold if $\phi_{j,i}\hat{\tau}_{j,i} \leq \hat{\pi}_j^{-1}\eta_i$. We consider the cases $j \in \dot{S}(i)$ and $j \in S(i) \setminus \dot{S}(i)$ separately. In the case $j \in \dot{S}(i)$, this is inequality immediate from (3.10a) in Assumption 3.1 (d) and Lemma 3.4. If $j \in S(i) \setminus \dot{S}(i)$, (3.10a) and Assumption 3.1 (b) give

$$\phi_{j,i}\hat{\tau}_{j,i}(\pi_j - \hat{\pi}_j) \leq \eta_{\tau,i}^\perp \leq \min_{j'}(\pi_{j'} - \hat{\pi}_{j'})\eta_i \leq (\pi_j - \hat{\pi}_j)\eta_i \leq (\pi_j - \hat{\pi}_j)\hat{\pi}_j^{-1}\eta_i.$$

In the last step we have used that $\hat{\pi}_j \in (0, 1]$ by (4.1). This finishes verifying (4.7).

Next, we expand (3.11a), obtaining

$$\begin{aligned} q_{j,i+2}(\bar{\gamma}_j) &= (2\mathbb{E}[\gamma_{j,i}\eta_i - \bar{\gamma}_j\phi_{j,i}\hat{\tau}_{j,i}|\mathcal{O}_i] + 2\alpha_i|\mathbb{E}[\gamma_{j,i}\eta_i - \bar{\gamma}_j\phi_{j,i}\hat{\tau}_{j,i}|\mathcal{O}_i]| - \delta\phi_{j,i})\chi_{S(i)}(j), \\ &= (2(\gamma_{j,i}\eta_i - \bar{\gamma}_j\phi_{j,i}\hat{\tau}_{j,i}) + 2\alpha_i|\gamma_{j,i}\eta_i - \bar{\gamma}_j\phi_{j,i}\hat{\tau}_{j,i}| - \delta\phi_{j,i})\chi_{S(i)}(j), \\ &\leq (2(1 + \alpha_i)\rho_j + 2(\bar{\gamma}_j\eta_i - \bar{\gamma}_j\phi_{j,i}\hat{\tau}_{j,i}) + 2\alpha_i|\bar{\gamma}_j\eta_i - \bar{\gamma}_j\phi_{j,i}\hat{\tau}_{j,i}| - \delta\phi_{j,i})\chi_{S(i)}(j). \end{aligned}$$

Since η_i and $\phi_{j,i}\tau_{j,i}$ are increasing, if also

$$(4.9) \quad 2(\bar{\gamma}_j\eta_i - \bar{\gamma}_j\phi_{j,i}\hat{\tau}_{j,i}) + 2\alpha_i|\bar{\gamma}_j\eta_i - \bar{\gamma}_j\phi_{j,i}\hat{\tau}_{j,i}| \leq \delta\phi_{j,i} \quad (j \in S(i)),$$

then

$$(4.10) \quad \mathbb{E}[q_{j,i+2}(\bar{\gamma}_j)] \leq 2(1 + \alpha_i)\rho_j.$$

Inserting α_i from (4.8), we see (4.9) to follow from (4.3d). Finally, (4.7) and (4.10) show that (3.12b) holds with $\rho_j = 0$. Thus Assumption 3.1 (e) holds. From Proposition 3.2 now

$$\delta_{j,i+2}^x(\bar{\gamma}_j) = 8(1 + \alpha_i)\rho_j C_x + 2(1 + \alpha_i^{-1})\rho_j C_x.$$

Clearly α_i defined in (4.8) is bounded above and below, so we obtain (4.3e). \square

4.2 THE PARAMETERS $\eta_{\tau,i}^\perp$ AND $\eta_{\sigma,i}^\perp$

We now want to satisfy [Assumption 3.1 \(c-i\)](#) for doubly-stochastic methods. As it turns out, the parameters $\eta_{\tau,i}^\perp$ and $\eta_{\sigma,i}^\perp$, do not have any effect on convergence rates. Here are a few options.

Lemma 4.2. *Assume [\(4.1\)](#) and that $i \mapsto \eta_i$ is non-decreasing with $\eta_i \in \mathcal{R}(\mathcal{O}_{i-1}; (0, \infty))$. Then [Assumption 3.1 \(b\)](#) & [\(c-i\)](#) hold and both $i \mapsto \eta_{\tau,i}^\perp$ and $i \mapsto \eta_{\sigma,i}^\perp$ are non-decreasing if either:*

(i) *(Constant rule) We take $\eta_{\tau,i}^\perp \equiv \eta_\tau^\perp$ and $\eta_{\sigma,i}^\perp \equiv \eta_\sigma^\perp$ for constant $\eta_\sigma^\perp, \eta_\tau^\perp > 0$ satisfying*

$$\eta_0 \cdot \min_j (\pi_j - \dot{\pi}_j) \geq \eta_\tau^\perp, \quad \text{and} \quad \eta_0 \cdot \min_\ell (v_\ell - \dot{v}_\ell) \geq \eta_\sigma^\perp.$$

(ii) *(Proportional rule) For some $\alpha \in (0, 1)$ we take $\eta_{\tau,i}^\perp := \eta_{\sigma,i}^\perp := \alpha \eta_i$ satisfying*

$$\min_j (\pi_j - \dot{\pi}_j) \geq \alpha, \quad \text{and} \quad \min_\ell (v_\ell - \dot{v}_\ell) \geq \alpha.$$

Proof. Clearly both rules satisfy [Assumption 3.1 \(b\)](#) & [\(c-i\)](#). That $i \mapsto \eta_{\tau,i}^\perp$ and $i \mapsto \eta_{\sigma,i}^\perp$ are non-decreasing and belong to $\mathcal{R}(\mathcal{O}_{i-1}; [0, \infty))$ is obvious. \square

4.3 WORST-CASE RULES FOR η_i

To verify [Assumption 3.1 \(a\)](#) we take deterministic worst-case bounds $w_j, w_{j,\ell} \geq 0$ such that

$$(4.11) \quad w_j := \max_\ell w_{\ell,j} \quad \text{and} \quad w_{\ell,j} \geq \dot{\pi}_j^{-1} \chi_{\dot{S}(j)}(j) + \dot{v}_\ell^{-1} \chi_{\dot{V}(i+1)}(\ell) \quad (i \in \mathbb{N}).$$

Since we assume iteration-independent probabilities [\(4.1\)](#), such bounds exist.

Lemma 4.3. *Suppose [Assumption 3.1 \(d\)](#) and [\(4.1\)](#) hold. With $(\kappa_1, \dots, \kappa_n) \in \mathcal{K}(K, \mathcal{P}, \mathcal{Q})$ take*

$$(4.12) \quad \eta_i := \min_{\ell=1, \dots, n} \sqrt{\frac{(1-\delta)\psi_{\ell,i+1}}{\kappa_\ell(w_{\ell,1}^2 \phi_{1,i}^{-1}, \dots, w_{\ell,m}^2 \phi_{m,i}^{-1})}} \quad (i \geq 0).$$

Then [Assumption 3.1 \(a\)](#) holds. Moreover, $\eta_i \in \mathcal{R}(\mathcal{O}_{i-1}; (0, \infty))$ provided $\psi_{i+1} \in \mathcal{R}(\mathcal{O}_{i-1}; (0, \infty))$.

Proof. Recalling the expression for $\lambda_{\ell,j,i}$ in [\(3.9\)](#), [Assumption 3.1 \(d\)](#), and [\(4.1\)](#) imply $\lambda_{\ell,j,i} \leq \eta_i w_{\ell,j}$ for $\ell \in \mathcal{V}(j)$. By the monotonicity of κ_ℓ (assumed in [Definition 2.2](#)), [Assumption 3.1 \(a\)](#) will therefore hold if

$$(4.13) \quad \psi_{\ell,i+1} \geq \frac{\eta_i^2}{1-\delta} \kappa_\ell(w_{\ell,1}^2 \phi_{1,i}^{-1}, \dots, w_{\ell,m}^2 \phi_{m,i}^{-1}).$$

This is verified by inserting η_i from [\(4.12\)](#). Clearly [\(4.12\)](#) also verifies $\eta_i \in \mathcal{R}(\mathcal{O}_{i-1}; (0, \infty))$ when $\psi_{i+1} \in \mathcal{R}(\mathcal{O}_{i-1}; (0, \infty))$. \square

The next lemma provides a choice of $\psi_{i+1} \in \mathcal{R}(\mathcal{O}_{i-1}; (0, \infty))$ that also satisfies [Assumption 3.1 \(f\)](#). The resulting η_i we express later in [\(4.14\)](#) to collect all rules in one place.

Lemma 4.4. Let $(\kappa_1, \dots, \kappa_n) \in \mathcal{K}(K, \mathcal{P}, \mathcal{Q})$ and take η_i according to (4.12) with $\psi_{\ell, i+1} = \eta_i^{2-1/p} \psi_{\ell, 0}$ for some $\psi_{\ell, 0} > 0$ and $p \in (0, 1]$. If $\phi_{j, i} \in \mathcal{R}(\mathcal{O}_{i-1}; (0, \infty))$, then $\eta_i, \psi_{i+1} \in \mathcal{R}(\mathcal{O}_{i-1}; (0, \infty))$. If, moreover, (4.3c) holds, then $\mathbb{E}[\eta_i] \geq c_\eta^p i^p$ and $\eta_i \geq b_\eta^p \min_j \phi_{j, i}^p$ for some constants $c_\eta, b_\eta > 0$ independent of p .

Proof. That $\eta_i, \psi_{i+1} \in \mathcal{R}(\mathcal{O}_{i-1}; (0, \infty))$ is clear from (4.12) and $\phi_{j, i} \in \mathcal{R}(\mathcal{O}_{i-1}; (0, \infty))$.

With $\underline{\psi}_0 := \min_{\ell=1, \dots, n} \psi_{\ell, 0}$, from (4.12) also

$$\eta_i^{1/p} \geq \frac{(1-\delta)\underline{\psi}_0}{\max_{\ell=1, \dots, n} \kappa_\ell (\mathbb{w}_{\ell, 1}^2 \phi_{1, i}^{-1}, \dots, \mathbb{w}_{\ell, m}^2 \phi_{m, i}^{-1})}.$$

Since $\hat{\mu}_{\ell, j, i} = 0$ for $\ell \notin \mathcal{V}(j)$, using Definition 2.2 (ii), we get

$$\eta_i^{1/p} \geq \frac{(1-\delta)\underline{\psi}_0}{\bar{\kappa} \sum_{j=1}^n \max_{\ell} \mathbb{w}_{\ell, j}^2 \phi_{j, i}^{-1}} \geq \frac{1}{\sum_{j=1}^n b_j^{-1} \phi_{j, i}^{-1}}$$

for $b_j := (1-\delta)\underline{\psi}_0 / (\bar{\kappa} \mathbb{w}_j^2)$. This shows $\eta_i \geq \min_j b_j^p \phi_{j, i}^p$. Since $x \mapsto 1/x$ and $x \mapsto x^q$ are convex on $[0, \infty)$ for $q \geq 1$, Jensen's inequality gives

$$\mathbb{E}[\eta_i] \geq \frac{1}{\mathbb{E}[(\sum_{j=1}^n b_j^{-1} \phi_{j, i}^{-1})^p]} \geq \frac{1}{(\sum_{j=1}^n b_j^{-1} \mathbb{E}[\phi_{j, i}^{-1}])^p}.$$

By an application of (4.3c) we obtain $\mathbb{E}[\eta_i] \geq c_\eta^p i^p$ for $c_\eta := 1 / \sum_{j=1}^m b_j^{-1} c_j$. \square

4.4 MIXED RATES UNDER PARTIAL STRONG CONVEXITY

We are finally ready to state our main result and algorithms. We recall that by Lemma 2.2, (PP) is equivalent to (2.19) under the structural conditions of Assumption 2.1. Dividing the updates of (2.19) into individual block updates, and taking the step length rules from Assumption 3.1 (d), we obtain the steps of the doubly-stochastic method Algorithm 1. If we perform full dual updates, i.e., force Assumption 3.1 (c-ii) and following Lemma 3.8 take $\mathring{V}(i+1) = \{1, \dots, n\}$ and $\mathring{S}(i) = S(i)$, we get the simpler steps of Algorithm 2. Regarding the updates of the remaining parameters that are not specified directly in the algorithm skeletons, we start with:

Theorem 4.5. Assume the block-separable structure (GF), writing $\gamma_j \geq 0$ for the factor of (strong) convexity of G_j . Let $\delta \in (0, 1)$ and $(\kappa_1, \dots, \kappa_n) \in \mathcal{K}(K, \mathcal{P}, \mathcal{Q})$ (see Definition 2.2). In Algorithm 1 or Algorithm 2, take

- (i) $\phi_{j, 0} > 0$ freely and $\phi_{j, i+1} := \phi_{j, i} + 2(\bar{\gamma}_j \eta_i + \rho_j)$ for some $\rho_j \geq 0$ and $\bar{\gamma}_j \in [0, \gamma_j]$ with $\rho_j + \bar{\gamma}_j > 0$.
- (ii) $\psi_{\ell, 0} > 0$ freely and $\psi_{\ell, i} := \psi_{\ell, 0} \eta_i^{2-1/p}$ for some fixed $p \in [1/2, 1]$.
- (iii) $\eta_{\tau, i}^\perp, \eta_{\sigma, i}^\perp > 0$ (in Algorithm 1) following Lemma 4.2, and, with \mathbb{w}_j given by (4.11),

$$(4.14) \quad \eta_i := \min_{\ell=1, \dots, n} \left(\frac{(1-\delta)\psi_{\ell, 0}}{\kappa_\ell (\mathbb{w}_{\ell, 1}^2 \phi_{1, i}^{-1}, \dots, \mathbb{w}_{\ell, m}^2 \phi_{m, i}^{-1})} \right)^p.$$

Let $\widehat{u} \in H^{-1}(0)$, i.e., solve (OC), and suppose the following hold:

- (A) $\sup_{j=1, \dots, m} \rho_j = 0$ or $\sup_{j=1, \dots, m; i \in \mathbb{N}} \|x_j^{i+1} - \widehat{x}_j\|^2 \leq C_x$ for a constant $C_x > 0$.
- (B) $p = \frac{1}{2}$ or both $\sup_{\ell=1, \dots, n; i \in \mathbb{N}} \|y_\ell^{i+1} - \widehat{y}_\ell\|^2 \leq C_y$ and $\bar{y}_{j^*} = 0$ for some $j^* \in \{1, \dots, m\}$.
- (C) With $\ell^*(j)$ and $\underline{\kappa}$ given by [Definition 2.2](#), for some $\bar{y}_j \in [\underline{\gamma}_j, \gamma_j]$ for all $j = 1, \dots, m$ we have the initialisation bound

$$(4.15) \quad \bar{y}_j = \underline{\gamma}_j = 0 \quad \text{or} \quad \frac{2\bar{y}_j \underline{\gamma}_j}{\bar{y}_j - \underline{\gamma}_j} \left(\frac{1 - \delta}{\underline{\kappa} w_j} \right)^p \leq \delta \psi_{\ell^*(j), 0}^{-p} \phi_{j, 0}^{1-p}.$$

Then

$$(4.16) \quad \sum_{j=1}^m \frac{\delta \bar{c}_j \bar{y}_j}{2} \mathbb{E} [\|x_j^N - \widehat{x}_j\|^2] + g_{p, N} \leq \frac{\|u^0 - \widehat{u}\|_{Z_0, M_0}^2 + 18C_x (\sum_{j=1}^m \rho_j) N + \sum_{\ell=1}^n \psi_{\ell, 0} (C_* N^{2p-1} + \delta_*)}{2N^{p+1}},$$

when $N \geq 4$ and the weighted gap on the ergodic variables,

$$g_{p, N} := \begin{cases} c_p \mathcal{G}(\bar{x}_N, \bar{y}_N), & \text{Algorithm 1, } \bar{y}_j \leq \gamma_j/2 \text{ for all } j, \\ c_{*, p} \mathcal{G}(\bar{x}_{*, N}, \bar{y}_{*, N}), & \text{Algorithm 2, } \bar{y}_j \leq \gamma_j/2 \text{ for all } j, \\ 0, & \text{otherwise.} \end{cases}$$

The constants $C_*, \delta_* \geq 0$ are zero if $p = 1/2$ while the constants $c_p, c_{*, p} > 0$.

Remark 4.6. If $p = 1/2$, (4.16) yields a mixed $O(1/N^{3/2}) + O(1/N^{1/2})$ convergence rate. If $p = 1$, we get a mixed $O(1/N^2) + O(1/N)$ convergence rate.

Remark 4.7. [Theorem 4.5](#) is valid (with suitable constants) for general primal update rules as long as (4.3) holds and $i \mapsto \phi_{j, i}$ is non-decreasing. This is the case for the deterministic rule of [Lemma 4.1](#). For the random rule of [Example 3.1](#), the rest of the conditions hold, but we have not been able to verify (4.3f). This has the implication that only the gap estimates hold.

Proof. We use [Proposition 3.2](#), so need to verify [Assumptions 2.1](#) and [3.1](#). First of all, (R) follows from the updates rules for the testing and step length parameters, that only depend on previous realisations of $S(i)$ and $V(i+1)$. The rest of the conditions of [Assumption 2.1](#) are clear from [Lemma 2.2](#), the derivation of [Algorithms 1](#) and [2](#) from (2.19), and the requisite nesting condition (V) within the algorithms themselves.

Regarding the requirements (a)–(f) of [Assumption 3.1](#), we proceed as follows:

- (a) The choice $\psi_{\ell, i+1} := \eta_i^{2-1/p} \psi_{\ell, 0}$ in (ii) shows that (4.14) is equivalent to the formula (4.12) for η_i . Thus [Lemma 4.3](#) verifies (a).
- (b) It is clear that $i \mapsto \phi_{j, i}$ and $i \mapsto \psi_{\ell, i}$ are non-decreasing. Therefore (4.12) shows that $i \mapsto \eta_i$ is non-decreasing. Moreover, [Lemma 4.4](#) verifies that $\eta_i \in \mathcal{R}(O_{i-1}; (0, \infty))$.

[Algorithm 2](#) by construction satisfies [Assumption 3.1 \(c-ii\)](#) and has both $\dot{V}(i+1) = \emptyset$ and $V(i+1) = \{1, \dots, n\}$. It therefore suffices to refer to [Lemma 3.8](#).

[Algorithm 1](#), by its own statement, satisfies (4.1). Therefore, [Lemma 4.2](#) shows [Assumption 3.1 \(b\) & \(c-i\)](#), and that also $i \mapsto \eta_{\tau, i}^\perp$ is non-decreasing.

Algorithm 1. Doubly-stochastic primal–dual method

Require: $K \in \mathcal{L}(X; Y)$, $G \in C(X)$, and $F^* \in C(Y)$ with the separable structures (GF).

Require: Rules for $\phi_{j,i}$, $\psi_{\ell,i+1}$, η_{i+1} , $\eta_{\tau,i+1}^\perp$, $\eta_{\sigma,i+1}^\perp > 0$ from [Theorem 4.5](#), [Corollary 4.8](#), or [4.9](#).

Require: Sampling patterns for $S(i)$, $\mathring{S}(i)$, $V(i+1)$, and $\mathring{V}(i+1)$, ($i \in \mathbb{N}$), subject to the nesting condition (\mathcal{V}) (p. 7) with iteration-independent probabilities (4.1); see [Section 3.6](#).

1: Choose initial iterates $x^0 \in X$ and $y^0 \in Y$.

2: Initialise $\tau_{j,-1}, \sigma_{\ell,0} := 0$, ($j = 1, \dots, m$; $\ell = 1, \dots, m$).

3: **for all** $i \geq 0$ **until** a stopping criterion is satisfied **do**

4: Sample $\mathring{S}(i) \subset S(i) \subset \{1, \dots, m\}$ and $\mathring{V}(i+1) \subset V(i+1) \subset \{1, \dots, n\}$.

5: For each $j \in \mathring{S}(i)$, compute

$$\tau_{j,i} := \frac{\eta_{i-\phi_{j,i-1}\tau_{j,i-1}\chi_{S(i-1)\setminus\mathring{S}(i-1)}(j)}}{\phi_{j,i}\pi_{j,i}}, \quad \text{and}$$

$$x_j^{i+1} := (I + \tau_{j,i}\partial G_j)^{-1} \left(x_j^i - \tau_{j,i} \sum_{\ell \in \mathcal{V}(j)} K_{\ell,j}^* y_\ell^i \right), \quad \text{where } K_{\ell,j} := Q_\ell K P_j.$$

6: For each $\ell \in \mathring{V}(i+1)$, compute

$$\sigma_{j,i+1} := \frac{\eta_{i-\psi_{j,i}\sigma_{j,i}\chi_{V(i)\setminus\mathring{V}(i)}(j)}}{\psi_{j,i+1}\hat{\nu}_{\ell,i+1}}, \quad \text{and}$$

$$y_\ell^{i+1} := (I + \sigma_{\ell,i+1}\partial F_\ell^*)^{-1} \left(y_\ell^i + \sigma_{\ell,i+1} \sum_{j \in \mathcal{V}^{-1}(\ell)} K_{\ell,j} x_j^i \right).$$

7: For each $j \in \mathring{S}(i)$ and $\ell \in \mathcal{V}(j)$, set

$$\tilde{w}_{\ell,j}^{i+1} := \theta_{\ell,j,i+1}(x_j^{i+1} - x_j^i) + x_j^{i+1} \quad \text{with} \quad \theta_{\ell,j,i+1} := \frac{\tau_{j,i}\phi_{j,i}}{\sigma_{\ell,i+1}\psi_{\ell,i+1}}.$$

8: For each $\ell \in \mathring{V}(i+1)$ and $j \in \mathcal{V}^{-1}(\ell)$, set

$$\tilde{v}_{\ell,j}^{i+1} := b_{\ell,j,i+1}(y_\ell^{i+1} - y_\ell^i) + y_\ell^i \quad \text{with} \quad b_{\ell,j,i+1} := \frac{\sigma_{\ell,i+1}\psi_{\ell,i+1}}{\tau_{j,i}\phi_{j,i}}.$$

9: For each $j \in S(i) \setminus \mathring{S}(i)$, compute

$$\tau_{j,i} := \frac{\eta_{\tau,i}^\perp}{\phi_{j,i}(\pi_{j,i} - \hat{\pi}_{j,i})}, \quad \text{and}$$

$$x_j^{i+1} := (I + \tau_{j,i}\partial G_j)^{-1} \left(x_j^i - \tau_{j,i} \sum_{\ell \in \mathcal{V}(j)} K_{\ell,j}^* \tilde{v}_{\ell,j}^{i+1} \right).$$

10: For each $\ell \in V(i+1) \setminus \mathring{V}(i+1)$ compute

$$\sigma_{j,i+1} := \frac{\eta_{\sigma,i}^\perp}{\psi_{j,i+1}(\nu_{\ell,i+1} - \hat{\nu}_{\ell,i+1})}, \quad \text{and}$$

$$y_\ell^{i+1} := (I + \sigma_{\ell,i+1}\partial F_\ell^*)^{-1} \left(y_\ell^i + \sigma_{\ell,i+1} \sum_{j \in \mathcal{V}^{-1}(\ell)} K_{\ell,j} \tilde{w}_{\ell,j}^{i+1} \right).$$

11: **end for**

(c) Proved together with (b) above.

(d) These choices are encoded into [Algorithm 1](#). For [Algorithm 2](#) we recall [Lemma 3.8](#).

(e) We use [Lemma 4.1](#). We have already showed [Assumption 3.1\(b\) & \(d\)](#). Moreover, the algorithms satisfy the iteration-independent probability assumption (4.1). By (A), either

Algorithm 2. Block-stochastic primal–dual method, primal randomisation only

Require: $K \in \mathcal{L}(X; Y)$, $G \in C(X)$, and $F^* \in C(Y)$ with the separable structures (GF).

Require: Rules for $\phi_{j,i}, \psi_{\ell,i+1}, \eta_{i+1} \in \mathcal{R}(\mathcal{O}_i; (0, \infty))$ from [Theorem 4.5](#), [Corollary 4.8](#), or [4.9](#).

Require: Iteration-independent (4.1) sampling pattern for the set $S(i)$, ($i \in \mathbb{N}$); see [Section 3.6](#).

- 1: Choose initial iterates $x^0 \in X$ and $y^0 \in Y$.
- 2: **for all** $i \geq 0$ **until** a stopping criterion is satisfied **do**
- 3: Sample $S(i) \subset \{1, \dots, m\}$.
- 4: For each $j \notin S(i)$, set $x_j^{i+1} := x_j^i$.
- 5: For each $j \in S(i)$, with $\tau_{j,i} := \eta_i \pi_{j,i}^{-1} \phi_{j,i}^{-1}$, compute

$$x_j^{i+1} := (I + \tau_{j,i} \partial G_j)^{-1} \left(x_j^i - \tau_{j,i} \sum_{\ell \in \mathcal{V}(j)} K_{\ell,j}^* y_\ell^i \right), \quad \text{where } K_{\ell,j} := Q_\ell K P_j.$$

- 6: For each $j \in S(i)$ set

$$\bar{x}_j^{i+1} := \theta_{j,i+1} (x_j^{i+1} - x_j^i) + x_j^{i+1} \quad \text{with } \theta_{j,i+1} := \frac{\eta_i}{\pi_{j,i} \eta_{i+1}}.$$

- 7: For each $\ell \in \{1, \dots, n\}$ using $\sigma_{\ell,i+1} := \eta_{i+1} \psi_{\ell,i+1}^{-1}$, compute

$$y_\ell^{i+1} := (I + \sigma_{\ell,i+1} \partial F_\ell^*)^{-1} \left(y_\ell^i + \sigma_{\ell,i+1} \sum_{j \in \mathcal{V}^{-1}(\ell)} K_{\ell,j} \bar{x}_j^{i+1} \right).$$

- 8: **end for**
-

$\sup_j \rho_j = 0$ or (3.12a) holds. We still need to satisfy (4.3d). Using [Definition 2.2 \(iii\)](#) in (4.14), we estimate

$$(4.19) \quad \eta_i \leq \left(\frac{(1-\delta) \psi_{\ell^*(j),0}}{\underline{\kappa} \mathbb{W}_j} \phi_{j,i} \right)^p.$$

By (C), therefore, either $\tilde{y}_j = \bar{y}_j = 0$, or $2\tilde{y}_j \bar{y}_j \eta_i \leq \delta (\tilde{y}_j - \bar{y}_j) \phi_{j,0}^{1-p} \phi_{j,i}^p$. By (i), $i \mapsto \phi_{j,i}$, so this gives (4.3d). [Lemma 4.1](#) now shows [Assumption 3.1 \(e\)](#).

- (f) If $p = 1/2$, by [Remark 4.6](#), $\psi_{\ell,i} \equiv \psi_{\ell,0}$. Therefore (3.13b) holds. If $p \neq 1/2$, the same remark and (A) guarantee (3.13a).

With [Assumptions 2.1](#) and [3.1](#) now verified, [Proposition 3.2](#) provides the estimate

$$(4.20) \quad \sum_{k=1}^m \frac{\delta}{2\mathbb{E}[\phi_{k,N}^{-1}]} \cdot \mathbb{E} [\|x_k^N - \hat{x}_k\|^2] + \tilde{g}_N \leq \frac{1}{2} \|u^0 - \hat{u}\|_{Z_0 M_0}^2 + \sum_{j=1}^m \frac{1}{2} d_{j,N}^x(\tilde{y}_j) + \sum_{\ell=1}^n \frac{1}{2} d_{\ell,N}^y,$$

where \tilde{g}_N , $d_{j,N}^x(\tilde{y}_j)$ and $d_{\ell,N}^y$ are given in (3.15). To obtain convergence rates, we still need to further analyse this estimate, mainly ζ_N and $\zeta_{*,N}$ within \tilde{g}_N .

We start with ζ_N and $\zeta_{*,N}$. By [Lemma 4.2](#) for [Algorithm 1](#) and directly by [Assumption 3.1 \(c-ii\)](#) for [Algorithm 2](#), $i \mapsto \eta_{\tau,i}^\perp$ is non-decreasing (as is $i \mapsto \eta_{\sigma,i}^\perp$). We recall the coupling variable $\bar{\eta}_i$ from (3.3). Observe that (4.3c) holds as we have verified the conditions of [Lemma 4.1](#) above.

By Corollary 3.5 and Lemma 4.4, therefore, in both cases, (3.3a) and (3.3b), for some constant $c_\eta > 0$,

$$\bar{\eta}_i = \mathbb{E}[\eta_i + \eta_{\tau,i}^\perp - \eta_{\tau,i-1}^\perp] \geq \mathbb{E}[\eta_i] \geq c_\eta^p i^p.$$

Thus we estimate ζ_N from (3.4) as

$$(4.21) \quad \begin{aligned} \zeta_N &= \sum_{i=0}^{N-1} \bar{\eta}_i \geq \sum_{i=0}^{N-1} \mathbb{E}[\eta_i] \geq c_\eta^p \sum_{i=0}^{N-1} i^p \geq c_\eta^p \int_0^{N-2} x^p dx \\ &\geq \frac{c_\eta^p}{p+1} (N-2)^{p+1} \geq \frac{c_\eta^p}{2^{p+1}(p+1)} N^{p+1} =: c_p N^{p+1} \quad (N \geq 4). \end{aligned}$$

Similarly, for some $c_{*,p} > 0$, the quantity $\zeta_{*,N}$ defined in (3.4) satisfies

$$(4.22) \quad \zeta_{*,N} \geq \sum_{i=1}^{N-1} \mathbb{E}[\eta_i] \geq \frac{c_\eta^p}{p+1} ((N-2)^{p+1} - 1) \geq c_{*,p} N^{p+1} \quad (N \geq 4).$$

If $p = 1/2$, (ii) clearly implies $d_{\ell,N}^y = \mathbb{E}[\psi_{\ell,N} - \psi_{\ell,0}] \equiv 0$. Therefore, we can take $C_*, \delta_* = 0$. Otherwise, since $0 \leq 2 - 1/p \leq 1$, the map $t \mapsto t^{2-1/p}$ is concave. Therefore, using (3.15), (ii), and Jensen's inequality, we deduce

$$d_{y,\ell}^N = \sum_{i=0}^{N-1} 9C_y \mathbb{E}[\psi_{\ell,i+2} - \psi_{\ell,i+1}] = 9C_y \psi_{\ell,0} (\mathbb{E}[\eta_{N+1}^{2-1/p}] - \mathbb{E}[\eta_1^{2-1/p}]) \leq 9C_y \psi_{\ell,0} \mathbb{E}[\eta_{N+1}]^{2-1/p}.$$

The condition (B) provides $j^* \in \{1, \dots, m\}$ with $\gamma_{j^*} = 0$, so that a referral to (4.3b) shows $\mathbb{E}[\phi_{j^*,N}] = \phi_{j^*,0} + 2N\rho_{j^*}$. By (4.19) for some $C_*, \delta_* > 0$ then

$$(4.23) \quad d_{y,\ell}^N \leq 9C_y \psi_{\ell,0} \left(\frac{(1-\delta)\psi_{\ell^*(j^*),0}}{\kappa_{\mathbb{W}_{j^*}}} \mathbb{E}[\phi_{j^*,i}] \right)^{2p-1} \leq \psi_{\ell,0} (C_* N^{2p-1} + \delta_*).$$

Finally, to estimate $d_{j,N}^x(\tilde{\gamma}_j)$, Lemma 4.4 shows $\eta_i \geq b_\eta^p \min_j \phi_{j,i}^p$, ($j = 1, \dots, m$). Thus (4.3f) and (4.3e) in Lemma 4.1 give $1/\mathbb{E}[\phi_{j,N}^{-1}] \geq \tilde{\gamma}_j \tilde{c}_j N^{p+1}$ for $N \geq 4$, and $d_{j,N}^x(\tilde{\gamma}_j) = 18\rho_j C_x N$. Now (4.16) is immediate by applying these estimates and (4.21)–(4.23) to (4.20). \square

4.5 UNACCELERATED ALGORITHM

If $\rho_j = 0$ and $\tilde{\gamma}_j = \tilde{\gamma}_j = 0$ for all $j = 1, \dots, m$, then $\phi_{j,i} \equiv \phi_{j,0}$. Consequently Lemma 4.3 gives $\eta_i \equiv \eta_0$. Recalling ζ_N from (3.4), we see that $\zeta_N = N\eta_0$. Likewise $\zeta_{*,N}$ from (3.4) satisfies $\zeta_{*,N} = (N-1)\eta_0$. Clearly also $d_{\ell,N}^y = 0$ and $d_{j,N}^x(\tilde{\gamma}_j) = 0$. Inserting this information into (4.20), we immediately obtain:

Corollary 4.8. *Assume the block-separable structure (GF). Let $\delta \in (0, 1)$ and $(\kappa_1, \dots, \kappa_n) \in \mathcal{K}(K, \mathcal{P}, Q)$. In Algorithm 1 or 2, take*

- (i) $\phi_{j,i} \equiv \phi_{j,0}$ for some fixed $\phi_{j,0} > 0$.
- (ii) $\psi_{\ell,i} \equiv \psi_{\ell,0}$ for some fixed $\psi_{\ell,0} > 0$.

(iii) $\eta_i \equiv \eta_0$ given by (4.12) and (in Algorithm 1) $\eta_{\tau,i}^\perp, \eta_{\sigma,i}^\perp > 0$ following Lemma 4.2.

Then

(I) The iterates of Algorithm 1 satisfy $\mathcal{G}(\tilde{x}_N, \tilde{y}_N) \leq C_0 \eta_0^{-1}/(2N)$, ($N \geq 1$).

(II) The iterates of Algorithm 2 satisfy $\mathcal{G}(\tilde{x}_{*,N}, \tilde{y}_{*,N}) \leq C_0/[2\eta_0(N-1)]$, ($N \geq 2$).

4.6 FULL PRIMAL STRONG CONVEXITY

If G is fully strongly convex, we can naturally derive an $O(1/N^2)$ algorithm.

Corollary 4.9. Assume the block-separable structure (GF), assuming each G_j , ($j = 1, \dots, m$), strongly convex with the corresponding factor $\gamma_j > 0$. Let $\delta \in (0, 1)$ and $(\kappa_1, \dots, \kappa_n) \in \mathcal{K}(K, \mathcal{P}, \mathcal{Q})$. In Algorithm 1 or Algorithm 2, take

(i) $\phi_{j,0} > 0$ freely and $\phi_{j,i+1} := \phi_{j,i}(1 + 2\tilde{\gamma}_j \tau_{j,i})$ for some fixed $\tilde{\gamma}_j \in (0, \gamma_j)$.

(ii) $\psi_{\ell,0} > 0$ freely and $\psi_{\ell,i} := \psi_{\ell,0}$.

(iii) η_i according to (4.12), and (in Algorithm 1) $\eta_{\tau,i}^\perp, \eta_{\sigma,i}^\perp > 0$ following Lemma 4.2.

Suppose the initialisation bound Theorem 4.5 (C) holds. Then

$$\sum_{j=1}^m \frac{\delta \tilde{c}_j \tilde{\gamma}_j}{2} \mathbb{E}[\|x_j^N - \hat{x}_j\|^2] + \tilde{g}_{1,N} \leq \frac{\|u^0 - \hat{u}\|_{Z_0 M_0}^2}{2N^2} \quad (N \geq 4)$$

for

$$\tilde{g}_{1,N} := \begin{cases} q_1 \mathcal{G}(\tilde{x}_N, \tilde{y}_N), & \text{Algorithm 1, } \tilde{\gamma}_j \leq \gamma_j/2 \text{ for all } j, \\ q_{*,1} \mathcal{G}(\tilde{x}_{*,N}, \tilde{y}_{*,N}), & \text{Algorithm 2, } \tilde{\gamma}_j \leq \gamma_j/2 \text{ for all } j, \\ 0, & \text{otherwise.} \end{cases}$$

The constants $\tilde{c}_j > 0$ are provided by Lemma 4.1 while $q_1, q_{*,1} > 0$.

Proof. We adapt the argumentation of Theorem 4.5 for the case $p = 1/2$. Indeed, with this choice, our present assumptions satisfy the conditions of that theorem hold:

(i) with $\rho_j = 0$ becomes the present one. Since we take $\tilde{\gamma}_j > 0$, $\rho_j + \tilde{\gamma}_j > 0$ as required.

(ii) reduces to the present one with $p = 1/2$.

(iii) becomes the present one since (4.14) with $p = 1/2$ equals (4.12).

(A) trivially holds since $\rho_j = 0$ for all $j = 1, \dots, m$.

(B) trivially holds when $p = 1/2$.

(C) we have assumed.

Since $C_*, \delta_* = 0$ when $p = 1/2$, the estimate (4.16) therefore holds with the right hand side $C_0/(2N^{1+1/2})$. We need to improve this to $C_0/(2N^2)$ by improving testing variable estimates.

Indeed, the update rule (4.2) now gives

$$\phi_{j,N} \geq \underline{\phi}_0 + \underline{\gamma} \sum_{i=0}^{N-1} \eta_i \geq \underline{\phi}_0 + \underline{\gamma} \sum_{i=0}^{N-1} \eta_i \quad \text{with} \quad \underline{\phi}_0 := \min_j \phi_{j,0} > 0.$$




Figure 1: Sample images for denoising, deblurring, and undimming experiments.

Lemma 4.4 shows $\eta_i^2 \geq \underline{b} \min_j \phi_{j,i}$ for some \underline{b} . Therefore $\eta_N^2 \geq \underline{b} \phi_0 + \underline{b} \gamma \sum_{i=0}^{N-1} \eta_i$. Otherwise written this says $\eta_N^2 \geq \tilde{\eta}_N^2$, where

$$\tilde{\eta}_N^2 = \underline{b} \phi_0 + \underline{b} \gamma \sum_{i=0}^{N-1} \tilde{\eta}_i = \tilde{\eta}_{N-1}^2 + c^2 \underline{\gamma} \tilde{\eta}_{N-1} = \tilde{\eta}_{N-1}^2 + \underline{b} \underline{\gamma} \tilde{\eta}_{N-1}^1.$$

This implies $\eta_i \geq \tilde{\eta}_i \geq c'_\eta i$ for some $c'_\eta > 0$; cf. the estimates for (2.3) in [6, 37]. Working through the final estimation stage of the proof of Theorem 4.5 with $p = 1/2$, we can now use use in (4.21) and (4.22) the estimate $\eta_i \geq c'_\eta i$ that would otherwise correspond to $p = 1$. In our final result, we write the constants c_p and $c_{*,p}$ from the proof as $q_1, q_{*,1} > 0$. \square

Remark 4.10 (Linear rates). *If both G and F^* are strongly convex, it is possible to derive linear rates. We refer to [35] for the single-block deterministic case.*

Remark 4.11 (Variance estimates). *Variance can be estimated following [35, Remark 3.4].*

5 NUMERICAL EXPERIENCE

We now apply several variants of the proposed algorithms to image processing problems. We consider discretisations, as our methods are formulated in Hilbert spaces, but the space of functions of bounded variation—where image processing problems are typically formulated—is only a Banach space. Our specific example problems will be TGV² denoising, TV deblurring, and TV undimming.

We present the corrupt and ground-truth images in Figure 1, with values in the range $[0, 255]$. We use the images both at the original resolution of $n_1 \times n_2 = 768 \times 512$, and scaled down to 192×128 pixels. To the noisy high-resolution test image in Figure 1b, we have added Gaussian noise with standard deviation 29.6 (12dB). In the downscaled image, this becomes 6.15 (25.7dB). The image in Figure 1c we have distorted with Gaussian blur of standard deviation 4. To avoid inverse crimes, we have added Gaussian noise of standard deviation 2.5. The dimmed image in Figure 1d, we have distorted by multiplying the image with a sinusoidal mask γ ; see Figure 1c. Again, we have added the small amount of noise.

Besides the unaccelerated PDHGM—our examples lack strong convexity for acceleration of basic methods—we evaluate our algorithms against the relaxed PDHGM of [7, 19]. In our precursor work [37], we have also evaluated these two algorithms against the mixed-rate method

Table 1: Algorithm variant name construction

Letter:	1st	2nd	3rd	4th
	Randomisation	ϕ rule	η and ψ rules	κ choice
A-	D: Deterministic P: Primal only B: Primal & Dual	R: Random, Lem. 3.1 D: Determ., Lem. 4.1 C: Constant	B: Bounded: $p = \frac{1}{2}$ I: Increasing: $p = 1$	O: Balanc., Ex. 2.2 M: Max., Ex. 2.1

of [8], and the adaptive PDHGM of [17]. To keep our tables and figures easily legible, we also do not include the algorithms of [37] in our evaluations. It is worth noting that even in the two-block case, the algorithms presented in this paper will not reduce to those of that paper: our rules for $\sigma_{\ell,i}$ are very different from the rules for the single σ_i therein.

We define abbreviations of our algorithm variants in Table 1. We do not report the results or apply all variants to all example problems, as this would not be informative. We demonstrate the performance of the stochastic variants on TGV² denoising only. This merely serves as an example, as our problems are not large enough to benefit from being split on a computer cluster, where the benefits of stochastic approaches would be apparent.

To rely on Theorem 4.5 for convergence, we still need to satisfy (3.13a) and (3.12a), or take $\rho_j = 0$. The bound C_y in Assumption 3.1 (f) is easily calculated, as in all of our example problems, the functional F^* will restrict the dual variable to lie in a ball of known size. The primal variable, on the other hand, is not explicitly bounded. It is however possible to prove data-based conservative bounds on the optimal solution, see, e.g., [36, Appendix A]. We can therefore add an artificial bound to the problem to force all iterates to be bounded, replacing G by $\tilde{G}(x) := G(x) + \delta_{B(0, C_x)}(x)$. In practise, to avoid figuring out the exact magnitude of C_x , we update it dynamically. This avoids the constraint from ever becoming active and affecting the algorithm at all. In [36] a “pseudo duality gap” based on this idea was introduced to avoid problems with numerically infinite duality gaps. We will also use them in our reporting: we take the bound C_x as the maximum over all iterations of all tested algorithms, and report the duality gap for the problem with \tilde{G} replacing G . We always report the pseudo-duality gap in decibels $10 \log_{10}(\text{gap}^2/\text{gap}_0^2)$ relative to the initial iterate.

In addition to the pseudo-duality gap, we report for each algorithm the distance to a target solution and function value. We report the distance in decibels $10 \log_{10}(\|v^i - \hat{v}\|^2/\|\hat{v}\|^2)$ and the primal objective value $\text{val}(x) := G(x) + F(Kx)$ relative to the target as $10 \log_{10}((\text{val}(x) - \text{val}(\hat{x}))^2/\text{val}(\hat{x})^2)$. The target solution \hat{x} we compute by taking one million iterations of the PDHGM. We performed our computations with Matlab+C-MEX on a MacBook Pro with 16GB RAM and a 2.8 GHz Intel Core i5 CPU. The initial iterates are $x^0 = 0$ and $y^0 = 0$.

5.1 TGV² DENOISING

In this problem, we write $x = (v, w)$ and $y = (\phi, \psi)$, where v is the image of interest, and take

$$G(x) = \frac{1}{2} \|f - v\|^2, \quad K = \begin{pmatrix} \nabla & -I \\ 0 & \mathcal{E} \end{pmatrix}, \quad \text{and} \quad F^*(y) = \delta_{B(0, \alpha)^{n_1 n_2}}(\phi) + \delta_{B(0, \beta)^{n_1 n_2}}(\psi).$$

Here $\alpha, \beta > 0$ are regularisation parameters, \mathcal{E} is the symmetrised gradient, and the balls are pixelwise Euclidean with the product Π over image pixels. Since there is no further spatial

Table 2: TGV² denoising performance: CPU time and number of iterations (at a resolution of 10) to reach given duality gap, distance to target, or primal objective value.

Method	low resolution						high resolution					
	gap ≤ -60 dB		tgt ≤ -60 dB		val ≤ -60 dB		gap ≤ -50 dB		tgt ≤ -50 dB		val ≤ -50 dB	
	iter	time	iter	time	iter	time	iter	time	iter	time	iter	time
PDHGM	30	0.21s	100	0.72s	110	0.79s	50	6.31s	870	111.83s	370	47.49s
Relax	20	0.20s	70	0.71s	70	0.71s	40	6.93s	580	102.89s	250	44.25s
A-DRIO	40	0.26s	230	1.55s	180	1.22s	70	9.17s	2750	365.52s	1050	139.48s
A-DRBO	80	0.54s	890	6.07s	500	3.41s	80	10.56s	860	114.81s	420	56.00s
A-DDIO	20	0.14s	50	0.36s	110	0.80s	60	7.37s	2140	267.29s	900	112.34s
A-DDBO	30	0.19s	50	0.32s	90	0.58s	60	7.85s	600	79.67s	340	45.09s

non-uniformity in this problem, it is natural to take as our projections $P_1x = v$, $P_2x = w$, $Q_1y = \phi$, and $Q_2y = \psi$. It is then not difficult to calculate the optimal κ_ℓ of Example 2.2, so we use only the ‘xxxO’ variants of the algorithms in Table 1.

As the regularisation parameters (β, α) , we choose $(4.4, 4)$ for the downscaled image. For the original image we scale these parameters by $(0.25^{-2}, 0.25^{-1})$ corresponding to the image downscaling factor [13]. Since G is not strongly convex with respect to w , we have $\tilde{\gamma}_2 = 0$. For v we take $\tilde{\gamma}_1 = 1/2$, corresponding to the gap versions of our convergence estimates.

We take $\delta = 0.01$, and parametrise the standard PDHGM with $\sigma_0 = 1.9/\|K\|$ and $\tau_0 \approx 0.52/\|K\|$ solved from $\tau_0\sigma_0 = (1 - \delta)\|K\|^2$. These are values that typically work well. For forward-differences discretisation of TGV² with cell width $h = 1$, we have $\|K\|^2 \leq 11.4$ [36]. For the ‘Relax’ method from [7], we use the same σ_0 and τ_0 , as well as the value 1.5 for the inertial ρ parameter. For the increasing- ψ ‘xxIx’ variants of our algorithms, we take $\rho_1 = \rho_2 = 5$, $\tau_{1,0} = \tau_0$, and $\tau_{2,0} = 3\tau_0$. For the bounded- ψ ‘xxBx’ variants we take $\rho_1 = \rho_2 = 5$, $\tau_{1,0} = \tau_0$, and $\tau_{2,0} = 8\tau_0$. For both methods we also take $\eta_0 = 1/\tau_{0,1}$. *These parametrisations force $\phi_{1,0} = 1/\tau_{1,0}^2$, and keep the initial step length $\tau_{1,0}$ for v consistent with the basic PDHGM.* This justifies our algorithm comparisons using just a single set of parameters. We plot the step length evolution for the A-DDBO variant in Figure 3a.

The results for deterministic variants of our algorithm are in Table 2 and Figure 2. We display the first 5000 iterations in a logarithmic fashion. To reduce computational overheads, we compute the reported quantities only every 10 iterations. To reduce the effects of other processes occasionally occupying the computer, the CPU times reported are the average `iteration_time = total_time/total_iterations`, excluding time spent initialising the algorithm.

Our first observation is that the variants ‘xDxx’ based on the deterministic ϕ rule perform better than the “random” rule ‘xRxx’. Presently, with no randomisation, the only difference is the value of $\tilde{\gamma}$. The value 0.0105 from the initialisation bound Theorem 4.5 (C) for $p = 1/2$ and the value 0.0090 for $p = 1$ appear to give better performance than the maximal value $\tilde{\gamma}_1 = 0.5$. Generally, the A-DDBO seems to have the best asymptotic performance, with A-DRBO close. A-DDIO has good initial performance, although especially on the higher resolution image, the PDHGM and ‘Relax’ perform initially the best. Overall, however, the question of the best performer seems to be a rather fair competition between ‘Relax’ and A-DDBO.




Figure 2: TGV² denoising, deterministic variants of our algorithms with pixelwise step lengths, 5000 iterations, high (hi-res) and low (lo-res) resolution images.

5.2 TGV² DENOISING WITH STOCHASTIC ALGORITHM VARIANTS

We also test stochastic variants of our algorithms based on the alternating sampling of [Example 3.3](#) with $M = 1$ and, when appropriate, [Example 3.4](#). We take all probabilities equal to 0.5, that is $\mathbb{p}_x = \tilde{\pi}_1 = \tilde{\pi}_2 = \tilde{v}_1 = \tilde{v}_2 = 0.5$. In the doubly-stochastic ‘Bxxx’ variants of the algorithms, we take $\eta_{\tau,i}^\perp = \eta_{\sigma,i}^\perp = 0.9 \cdot 0.5\eta_i$ following the proportional rule [Lemma 4.2\(ii\)](#).

The results are in [Figure 4](#). To conserve space, we have only included a few descriptive algorithm variants. On the x axis, to better describe to the amount of actual work performed by the stochastic methods, the “iteration” count refers to the *expected* number of full primal–dual updates. For all the displayed stochastic variants, with the present choice of probabilities, the expected number of full updates in each iteration is 0.75.

We run each algorithm 50 times, and plot for each iteration the 90% confidence interval according to Student’s t -distribution. Towards the 5000th iteration, these generally become very narrow, indicating reliability of the random method. Overall, the full-dual-update ‘Pxxx’ variants perform better than the doubly-stochastic ‘Bxxx’ variants. In particular, A-PDBO has performance comparable to or even better than the PDHGM.

5.3 TV DEBLURRING

We want to remove the blur in [Figure 1c](#). We do this by taking

$$G(x) = \frac{1}{2} \|f - \mathcal{F}^*(a\mathcal{F}x)\|^2, \quad K = \nabla, \quad \text{and} \quad F^*(y) = \delta_{B(0,\alpha)^{n_1 n_2}}(y),$$




Figure 3: Step length evolution (logarithmic from initialisation). A-DDBO TGV² denoising and A-DDIM TV deblurring. The τ plots of the latter are images in the Fourier domain, lighter colour means smaller value of τ relative to initialisation (for that specific Fourier component). Note that the images depict logarithm change, not absolute values.




Figure 4: TGV² denoising, stochastic variants of our algorithms: 5000 iterations, low resolution images. Iteration number scaled by the fraction of blocks updated on average. For each iteration, 90% confidence interval according to the t -distribution over 50 random runs.




Figure 5: TV deblurring, deterministic variants of our algorithms with pixelwise step lengths, first 5000 iterations, high (hi-res) and low (lo-res) resolution images.

where the balls are again pixelwise Euclidean, and \mathcal{F} the discrete Fourier transform. The factors $a = (a_1, \dots, a_m)$ model the blurring operation in Fourier basis.

We take $\alpha = 2.55$ for the high resolution image and scale this to $\alpha = 2.55 * 0.15$ for the low resolution image. We parametrise the PDHGM and ‘Relax’ algorithms exactly as for TGV² denoising above, taking into account the estimate $8 \geq \|K\|^2$ [5]. We take $Q_1 = I$ and P_j as the projection to the j :th Fourier component so $m = n_1 n_2$ and $n = 1$. Thus *each primal Fourier component has its own step length parameter*. We initialise the latter as $\tau_{j,0} = \tau_0 / (\lambda + (1 - \lambda)\gamma_j)$, where the componentwise factor of strong convexity $\gamma_j = |a_j|^2$. For the bounded- ψ ‘xxBx’ algorithm variants we take $\lambda = 0.01$, and for the increasing- ψ ‘xxIx’ variants $\lambda = 0.1$. We illustrate the step length evolution of the variant A-DDIM in Figure 3.

We only experiment with deterministic algorithms, as we do not expect small-scale randomisation to be beneficial. We also use the maximal κ ‘xxxM’ variants, as a more optimal κ would be difficult to compute. The results are in Table 3 and Figure 5. Similarly to A-DDBO in our TGV² denoising experiments, A-DDBM performs reliably well, indeed better than the PDHGM or ‘Relax’. However, in many cases, A-DRBM and A-DDIM are even faster.

5.4 TV UNDIMMING

We take K and F^* as for TV deblurring, but $G(u) := \frac{1}{2} \|f - \gamma \cdot u\|^2$ for the sinusoidal dimming mask $\gamma : \Omega \rightarrow \mathbb{R}$. Our experimental setup is also nearly the same as TV deblurring, with the natural difference that the projection P_j are no longer to the Fourier basis, but to individual image pixels. The results are in Figure 6, and Table 4. They tell roughly the same story as TV deblurring, with A-DDBM performing well and reliably.

Table 3: TV deblurring performance: CPU time and number of iterations (at a resolution of 10) to reach given duality gap, distance to target, or primal objective value.

Method	low resolution						high resolution					
	gap ≤ -60 dB		tgt ≤ -60 dB		val ≤ -60 dB		gap ≤ -50 dB		tgt ≤ -40 dB		val ≤ -40 dB	
	iter	time	iter	time	iter	time	iter	time	iter	time	iter	time
PDHGM	30	0.18s	330	2.05s	70	0.43s	60	5.04s	330	28.12s	110	9.31s
Relax	20	0.11s	220	1.30s	50	0.29s	50	4.32s	220	19.30s	90	7.84s
A-DRIM	20	0.14s	280	2.08s	80	0.59s	30	3.27s	280	31.41s	320	35.92s
A-DRBM	20	0.14s	490	3.58s	90	0.65s	60	6.48s	240	26.27s	220	24.07s
A-DDIM	20	0.14s	170	1.25s	70	0.51s	30	3.17s	260	28.35s	230	25.06s
A-DDBM	20	0.15s	180	1.37s	60	0.45s	50	5.56s	230	25.98s	150	16.90s




Figure 6: TV undimming, deterministic variants of our algorithms with pixelwise step lengths, 5000 iterations, high (hi-res) and low (lo-res) resolution images.

CONCLUSIONS

We have derived several accelerated block-proximal primal–dual methods, both stochastic and deterministic. We have concentrated on applying them deterministically, taking advantage of blockwise—indeed pixelwise—factors of strong convexity, to obtain improved performance compared to standard methods. In future work, it will be interesting to evaluate the methods on real large scale problems to other state-of-the-art stochastic optimisation methods. Moreover, interesting questions include heuristics and other mechanisms for optimal initialisation of the pixelwise parameters, as well as combination with over-relaxation and inertial schemes, such as the extensions of the PDHGM considered in [10, 18, 34, 38].

Table 4: TV undimming performance: CPU time and number of iterations (at a resolution of 10) to reach given duality gap, distance to target, or primal objective value.

Method	low resolution						high resolution					
	gap ≤ -80 dB		tgt ≤ -60 dB		val ≤ -60 dB		gap ≤ -80 dB		tgt ≤ -60 dB		val ≤ -60 dB	
	iter	time	iter	time	iter	time	iter	time	iter	time	iter	time
PDHGM	70	0.18s	200	0.51s	120	0.30s	100	3.41s	300	10.31s	210	7.21s
Relax	50	0.16s	130	0.41s	80	0.25s	70	3.03s	200	8.73s	140	6.10s
A-DRIM	30	0.10s	160	0.57s	80	0.28s	80	3.52s	760	33.82s	640	28.48s
A-DRBM	20	0.05s	170	0.47s	60	0.16s	90	3.95s	370	16.39s	380	16.84s
A-DDIM	30	0.08s	110	0.30s	60	0.16s	70	3.05s	580	25.57s	430	18.94s
A-DDBM	20	0.05s	70	0.18s	40	0.10s	60	2.63s	230	10.22s	200	8.88s

ACKNOWLEDGEMENTS

The author would like to thank Peter Richtárik and Olivier Fercoq for several fruitful discussions, and for introducing him to stochastic optimisation. Moreover, the support of the EPSRC grant EP/M00483X/1 “Efficient computational tools for inverse imaging problems” is acknowledged during the initial two months of the research.

A DATA STATEMENT FOR THE EPSRC

Implementations of the algorithms described in the paper, and relevant boilerplate codes, are available on Zenodo at doi:[10.5281/zenodo.1042419](https://doi.org/10.5281/zenodo.1042419). The sample photo, also included in the archive, is from the free Kodak image suite, at the time of writing at <http://rok.us/graphics/kodak/>.

REFERENCES

- [1] BECK & TEOULLE, A Fast Iterative Shrinkage-Thresholding Algorithm for Linear Inverse Problems, *SIAM Journal on Imaging Sciences* 2 (2009), 183–202, DOI: [10.1137/080716542](https://doi.org/10.1137/080716542).
- [2] BERTSEKAS, Incremental aggregated proximal and augmented Lagrangian algorithms, 2015, ARXIV: [1509.09257](https://arxiv.org/abs/1509.09257).
- [3] BIANCHI, HACHEM & LUTZELER, A Stochastic Coordinate Descent Primal-Dual Algorithm and Applications to Large-Scale Composite Optimization, ARXIV: [1407.0898](https://arxiv.org/abs/1407.0898).
- [4] BOLTE, SABACH & TEOULLE, Proximal alternating linearized minimization for nonconvex and nonsmooth problems, *Mathematical Programming* 146 (2013), 459–494, DOI: [10.1007/s10107-013-0701-9](https://doi.org/10.1007/s10107-013-0701-9).
- [5] CHAMBOLLE, An algorithm for mean curvature motion, *Interfaces and Free Boundaries* 6 (2004), 195.
- [6] CHAMBOLLE & POCK, A first-order primal-dual algorithm for convex problems with applications to imaging, *Journal of Mathematical Imaging and Vision* 40 (2011), 120–145, DOI: [10.1007/s10851-010-0251-1](https://doi.org/10.1007/s10851-010-0251-1).

- [7] CHAMBOLLE & POCK, On the ergodic convergence rates of a first-order primal–dual algorithm, *Mathematical Programming* (2015), 1–35, DOI: [10.1007/s10107-015-0957-3](https://doi.org/10.1007/s10107-015-0957-3).
- [8] CHEN, LAN & OUYANG, Optimal Primal-Dual Methods for a Class of Saddle Point Problems, *SIAM Journal on Optimization* 24 (2014), 1779–1814, DOI: [10.1137/130919362](https://doi.org/10.1137/130919362).
- [9] COMBETTES & PESQUET, Stochastic forward-backward and primal-dual approximation algorithms with application to online image restoration, 2016, ARXIV: [1602.08021](https://arxiv.org/abs/1602.08021).
- [10] CONDAT, A Primal–Dual Splitting Method for Convex Optimization Involving Lipschitzian, Proxiable and Linear Composite Terms, *Journal of Optimization Theory and Applications* 158 (2013), 460–479, DOI: [10.1007/s10957-012-0245-9](https://doi.org/10.1007/s10957-012-0245-9).
- [11] CSIBA, QU & RICHTÁRIK, Stochastic Dual Coordinate Ascent with Adaptive Probabilities, ARXIV: [1502.08053](https://arxiv.org/abs/1502.08053).
- [12] DAUBECHIES, DEFRISE & DE MOL, An iterative thresholding algorithm for linear inverse problems with a sparsity constraint, *Communications on Pure and Applied Mathematics* 57 (2004), 1413–1457, DOI: [10.1002/cpa.20042](https://doi.org/10.1002/cpa.20042).
- [13] DE LOS REYES, SCHÖNLIEB & VALKONEN, Bilevel parameter learning for higher-order total variation regularisation models, *Journal of Mathematical Imaging and Vision* 57 (2017), 1–25, DOI: [10.1007/s10851-016-0662-8](https://doi.org/10.1007/s10851-016-0662-8), ARXIV: [1508.07243](https://arxiv.org/abs/1508.07243),
- [14] ESSER, ZHANG & CHAN, A general framework for a class of first order primal-dual algorithms for convex optimization in imaging science, *SIAM Journal on Imaging Sciences* 3 (2010), 1015–1046, DOI: [10.1137/09076934X](https://doi.org/10.1137/09076934X).
- [15] FERCOQ & BIANCHI, A Coordinate Descent Primal-Dual Algorithm with Large Step Size and Possibly Non Separable Functions, 2015, ARXIV: [1508.04625](https://arxiv.org/abs/1508.04625).
- [16] FERCOQ & RICHTÁRIK, Optimization in High Dimensions via Accelerated, Parallel, and Proximal Coordinate Descent, *SIAM Review* 58 (2016), 739–771, DOI: [10.1137/16M1085905](https://doi.org/10.1137/16M1085905).
- [17] GOLDSTEIN, LI & YUAN, Adaptive Primal-Dual Splitting Methods for Statistical Learning and Image Processing, *Advances in Neural Information Processing Systems* 28 (2015), ed. by CORTES et al., 2080–2088.
- [18] HE, YOU & YUAN, On the Convergence of Primal-Dual Hybrid Gradient Algorithm, *SIAM Journal on Imaging Sciences* 7 (2014), 2526–2537, DOI: [10.1137/140963467](https://doi.org/10.1137/140963467).
- [19] HE & YUAN, Convergence Analysis of Primal-Dual Algorithms for a Saddle-Point Problem: From Contraction Perspective, *SIAM Journal on Imaging Sciences* 5 (2012), 119–149, DOI: [10.1137/100814494](https://doi.org/10.1137/100814494).
- [20] HE & YUAN, Block-wise alternating direction method of multipliers for multiple-block convex programming and beyond, *SMAI Journal of Computational Mathematics* 1 (2015), 145–174, DOI: [10.5802/smai-jcm.6](https://doi.org/10.5802/smai-jcm.6).
- [21] MÖLLENHOFF, STREKALOVSKIY, MOELLER & CREMERS, The primal-dual hybrid gradient method for semiconvex splittings, *SIAM Journal on Imaging Sciences* 8 (2015), 827–857, DOI: [10.1137/140976601](https://doi.org/10.1137/140976601).

- [22] NESTEROV, Efficiency of Coordinate Descent Methods on Huge-Scale Optimization Problems, *SIAM Journal on Optimization* 22 (2012), 341–362, DOI: [10.1137/100802001](https://doi.org/10.1137/100802001).
- [23] OCHS, CHEN, BROX & POCK, iPiano: inertial proximal algorithm for nonconvex optimization, *SIAM Journal on Imaging Sciences* 7 (2014), 1388–1419, DOI: [10.1137/130942954](https://doi.org/10.1137/130942954).
- [24] PENG, XU, YAN & YIN, ARock: an Algorithmic Framework for Asynchronous Parallel Coordinate Updates, CAM Report 15-37, UCLA, 2015, URL: <ftp://ftp.math.ucla.edu/pub/camreport/cam15-37.pdf>.
- [25] PENG et al., Coordinate Friendly Structures, Algorithms and Applications, 2016, ARXIV: [1601.00863](https://arxiv.org/abs/1601.00863).
- [26] PESQUET & REPETTI, A class of randomized primal-dual algorithms for distributed optimization, 2014, ARXIV: [1406.6404](https://arxiv.org/abs/1406.6404).
- [27] POCK, CREMERS, BISCHOF & CHAMBOLLE, An algorithm for minimizing the Mumford-Shah functional, in: *12th IEEE Conference on Computer Vision*, 2009, 1133–1140, DOI: [10.1109/ICCV.2009.5459348](https://doi.org/10.1109/ICCV.2009.5459348).
- [28] QU, RICHTÁRIK & ZHANG, Randomized Dual Coordinate Ascent with Arbitrary Sampling, 2014, ARXIV: [1411.5873](https://arxiv.org/abs/1411.5873).
- [29] RICHTÁRIK & TAKÁČ, Parallel coordinate descent methods for big data optimization, *Mathematical Programming* (2015), 1–52, DOI: [10.1007/s10107-015-0901-6](https://doi.org/10.1007/s10107-015-0901-6).
- [30] RICHTÁRIK & TAKÁČ, Distributed coordinate descent method for learning with big data, 2013, ARXIV: [1310.2059](https://arxiv.org/abs/1310.2059).
- [31] SHALEV-SHWARTZ & ZHANG, Accelerated proximal stochastic dual coordinate ascent for regularized loss minimization, *Mathematical Programming* 155 (2014), 105–145, DOI: [10.1007/s10107-014-0839-0](https://doi.org/10.1007/s10107-014-0839-0).
- [32] SHIR ĀEV, *Probability*, Springer, 1996.
- [33] SUZUKI, Stochastic Dual Coordinate Ascent with Alternating Direction Multiplier Method, 2013, ARXIV: [1311.0622](https://arxiv.org/abs/1311.0622).
- [34] VALKONEN, Inertial, corrected, primal–dual proximal splitting, 2018, ARXIV: [1804.08736](https://arxiv.org/abs/1804.08736), URL: tuomov.iki.fi/m/inertia.pdf.
- [35] VALKONEN, Testing and non-linear preconditioning of the proximal point method, *Applied Mathematics and Optimization* (2018), DOI: [10.1007/s00245-018-9541-6](https://doi.org/10.1007/s00245-018-9541-6), ARXIV: [1703.05705](https://arxiv.org/abs/1703.05705),
- [36] VALKONEN, BREDIES & KNOLL, Total generalised variation in diffusion tensor imaging, *SIAM Journal on Imaging Sciences* 6 (2013), 487–525, DOI: [10.1137/120867172](https://doi.org/10.1137/120867172),
- [37] VALKONEN & POCK, Acceleration of the PDHGM on partially strongly convex functions, *Journal of Mathematical Imaging and Vision* 59 (2017), 394–414, DOI: [10.1007/s10851-016-0692-2](https://doi.org/10.1007/s10851-016-0692-2), ARXIV: [1511.06566](https://arxiv.org/abs/1511.06566),
- [38] VŮ, A splitting algorithm for dual monotone inclusions involving cocoercive operators, *Advances in Computational Mathematics* 38 (2013), 667–681, DOI: [10.1007/s10444-011-9254-8](https://doi.org/10.1007/s10444-011-9254-8).

- [39] WRIGHT, Coordinate descent algorithms, *Mathematical Programming* 151 (2015), 3–34, DOI: [10.1007/s10107-015-0892-3](https://doi.org/10.1007/s10107-015-0892-3).
- [40] YU, LIN & YANG, Doubly Stochastic Primal-Dual Coordinate Method for Empirical Risk Minimization and Bilinear Saddle-Point Problem, 2015, ARXIV: [1508.03390](https://arxiv.org/abs/1508.03390).
- [41] ZHANG & XIAO, Stochastic primal-dual coordinate method for regularized empirical risk minimization, 2014, ARXIV: [1409.3257](https://arxiv.org/abs/1409.3257).
- [42] ZHAO & ZHANG, Stochastic optimization with importance sampling, 2014, ARXIV: [1401.2753](https://arxiv.org/abs/1401.2753).
- [43] ZHU & CHAN, An efficient primal-dual hybrid gradient algorithm for total variation image restoration, CAM Report 08-34, UCLA, 2008, URL: <ftp://ftp.math.ucla.edu/pub/camreport/camo8-34.pdf>.